

研究業績目録

原著論文

1. Buckle T, Van den Berg N.S., Kuil J, Bunschoten A, Oldenburg J, Borowsky A. D., Wesseling J, Masada R, Oishi S, Fujii N, Van Leeuwen F.B. Non-invasive longitudinal imaging of tumor progression using an 111Indium labeled CXCR4 peptide antagonist.
J. Nucl. Med. Mol. Imaging in press.
2. Suzuki Y, Naoe S, Oishi S, Fujii N, Ohno H. Gold-catalyzed three-component annulation: efficient synthesis of highly functionalized dihydropyrazoles from alkynes, hydrazines, and aldehydes or ketones.
Org. Lett. in press.
3. Inoue N, Sasagawa K, Ikai K, Sasaki Y, Tomikawa J, Oishi S, Fujii N, Uenoyama Y, Ohmori Y, Yamamoto N, Hondo E, Maeda K, Tsukamura H. Kisspeptin neurons mediate reflex ovulation in the musk shrew (*Suncus murinus*).
Proc Natl Acad Sci USA. **108**(42) 17527-17532. (2011)
4. Hirano K, Inaba Y, Takasu K, Oishi S, Takemoto Y, Fujii N, Ohno H. Gold(I)-catalyzed polycyclizations of polyenyne-type anilines based on hydroamination and consecutive hydroarylation cascade.
J. Org. Chem. **76**(21) 9068-9080. (2011)
5. Xu C, Liu J, Chen L, Liang S, Fujii N, Tamamura H, Xiong H. HIV-1 gp120 enhances outward potassium current via CXCR4 and cAMP-dependent protein kinase a signaling in cultured rat microglia.
Glia **59**(6) 997-1007. (2011)
6. Ohno H, Yamamoto M, Iuchi M, Fujii N, Tanaka T. Palladium-catalyzed construction of polycyclic heterocycles by an alkyne insertion and direct arylation cascade.
Synthesis (16) 2567-2578 (2011).
7. Kinoshita T, Sekiguchi Y, Fukada H, Nakaniwa T, Tada T, Nakamura S, Kitaura K, Ohno H, Suzuki Y, Hirasawa A, Nakanishi I, Tsujimoto G. A detailed thermodynamic profile of cyclopentyl and isopropyl derivatives binding to CK2 kinase.
Mol. Cell. Biochem. **356**(1-2) 97-105. (2011)

8. Yoshimitsu Y, Oishi S, Miyagaki J, Inuki S, Ohno H, Fujii N.
Pachastrissamine (jaspine B) and its stereoisomers inhibit sphingosine kinases and atypical protein kinase C.
Bioorg. Med. Chem. **19**(18) 5402-5408. (2011)
9. Nishizawa K, Nishiyama H, Matsui Y, Kobayashi T, Kotani H, Masutani H, Oishi S, Saito R, Toda Y, Fujii N, Yodoi J, Ogawa O.
Thioredoxin interacting protein suppresses bladder carcinogenesis.
Carcinogenesis in press.
10. Izumi K, Watanabe K, Oishi S, Fujii N, Matsuoka M, Sarafianosc SG, Kodama E.
Potent anti-HIV-1 activity of N-HR-derived peptides including a deep pocket-forming region without antagonistic effect on T-20.
Antivir. Chem. Chemother. **22**(1) 51-55. (2011)
11. Takeuchi T, Oishi S, Watanabe T, Ohno H, Sawada J, Matsuno K, Asai A, Asada N, Kitaura K, Fujii N.
Structure-activity relationships of carboline and carbazole derivatives as a novel class of ATP-competitive kinesin spindle protein inhibitors.
J. Med. Chem. **54**(13) 4839-4846. (2011)
12. Iwata A, Inuki S, Oishi S, Fujii N, Ohno H.
Formal total synthesis of (+)-Lysergic acid via zinc(II)-mediated regioselective ring-opening reduction of 2-alkynyl-3-indolyloxirane.
J. Org. Chem. **76**(13) 5506-5512. (2011)
13. Tokimizu Y, Ohta Y, Chiba H, Oishi S, Fujii N, Ohno H.
Direct synthesis of highly fused perimidines by copper(I)-catalyzed hydroamination of 2-ethynylbenzaldehydes.
Tetrahedron **67**(29) 5168-5175. (2011)
14. Kuil J, Buckle T, Yuan H, van den Berg N. S, Oishi S, Fujii N, Josephson L, van Leeuwen F. W. B.
Synthesis and evaluation of a bimodal CXCR4 antagonistic peptide.
Bioconjug. Chem. **22**(5) 859-864. (2011)
15. Xu C, Liu J, Chen L, Liang S, Fujii N, Tamamura H, Xiong H.
HIV-1 gp120 enhances outward potassium current via CXCR4 and cAMP-dependent protein kinase a signaling in cultured rat microglia.
Glia. **59**(6) 997-1007. (2011)
16. Yamada M, Kubo H, Kobayashi S, Ishizawa K, He M, Suzuki T, Fujino N, Kunishima H, Hatta M, Nishimaki K, Aoyagi T, Tokuda K, Kitagawa M,

- Yano H, Tamamura H, Fujii N, Kaku M.
The increase in surface CXCR4 expression on lung extravascular neutrophils and its effects on neutrophils during endotoxin-induced lung injury.
Cell. Mol. Immunol. **8**(4) 305-314. (2011)
17. Masuda R., Oishi S., Ohno H., Kimura H., Saji H., Fujii, N.
Concise site-specific synthesis of DTPA-peptide conjugates: application to imaging probes for the chemokine receptor CXCR4.
Bioorg. Med. Chem. **19**(10) 3216-3220. (2011)
18. Inokuchi E., Oishi S., Kubo T., Ohno H., Shimura K., Matsuoka M., Fujii, N.
Potent CXCR4 antagonists containing amidine-type peptide bond isosteres.
ACS Med. Chem. Lett. **2**(6) 477-480. (2011)
19. Xing N. D., Ding S. T., Saito R., Nishizawa K., Kobayashi T., Inoue T., Oishi S., Fujii N., Lv J. J., Ogawa O., Nishiyama, H.
A potent chemotherapeutic strategy in prostate cancer: S-(methoxytrityl)-L-cysteine, a novel Eg5 inhibitor.
Asian J. Androl. **13**(2) 236-241. (2011)
20. Inokuchi E., Yamada A., Hozumi K., Tomita K., Oishi S., Ohno H., Nomizu M., Fujii N.
Design and synthesis of amidine-type peptide bond isostere: application of nitrile oxide derivatives as active ester equivalents to peptide and peptidomimetics synthesis.
Org. Biomol. Chem. **9**(9) 3421-3427. (2011)
21. Dar A., Schajnovitz A., Lapid K., Kalinkovich A., Itkin T., Ludin A., Kao W., Battista M., Tesio M., Kollet O., Cohen N. N., Margalit R., Buss E. C., Baleux F., Oishi S., Fujii N., Larochelle A., Dunbar C. E., Broxmeyer H. E., Frenette P. S., Lapidot, T.
Rapid mobilization of hematopoietic progenitors by AMD3100 and catecholamines is mediated by CXCR4-dependent SDF-1 release from bone marrow stromal cells.
Leukemia. **25**(8) 1286-1296. (2011)
22. Kang HS., Baba T., Mandai M., Matsumura N., Hamanishi J., Kharma B., Kondoh E., Yoshioka Y., Oishi S., Fujii N., Murphy SK., Konishi I.
GPR54 is a target for suppression of metastasis in endometrial cancer.
Mol Cancer Ther. **10**(4) 580-590. (2011)
23. Inuki S., Iwata A., Oishi S., Fujii N., Ohno H.
Enantioselective total synthesis of (+)-Lysergic acid, (+)-Lysergol, and

- (+)-Isolysergol by palladium-catalyzed domino cyclization of allenes bearing amino and bromoindolyl groups.
J. Org. Chem. **76**(7) 2072-2083. (2011)
24. Hirano K., Inaba Y., Takahashi N., Shimano M., Oishi S., Fujii N., Ohno H. Direct synthesis of fused-indoles by gold-catalyzed cascade cyclization of diynes.
J. Org. Chem. **76**(5) 1212-1227. (2011)
25. Noda M., Omatsu Y., Sugiyama T., Oishi S., Fujii N., Nagasawa T. CXCL12-CXCR4 chemokine signaling is essential for NK cell development in adult mice.
Blood **117**(2) 451-458. (2011)
26. Oishi S., Misu R., Setsuda S., Masuda R., Ohno H., Naniwa Y., Ieda N., Inoue N., Ohkura S., Uenoyama Y., Tsukamura H., Maeda K., Hirasawa A., Tsujimoto G., Fujii N. Activation of neuropeptide FF receptors by kisspeptin receptor ligands.
ACS Med. Chem. Lett. **2**(1) 53-57. (2011)
27. Gravel S., Malouf C., Boulais P.E., Berchiche Y.A., Oishi S., Fujii N., Leduc R., Sinnott D., Heveker N. The peptidomimetic CXCR4 antagonist TC14012 recruits β -arrestin to CXCR7 roles of receptor domains.
J. Biol. Chem. **285**(49) 37939-37943. (2010)
28. Shimura K., Nameki D., Kajiwara K., Watanabe K., Sakagami Y., Oishi S., Fujii N., Matsuoka M., Sarafianos S.G., Kodama E. Resistance profiles of novel electrostatically constrained HIV-1 fusion inhibitors.
J. Biol. Chem. **285**(50) 39471-39480. (2010)
29. Martin S.K., Diamond P., Williams S.A., To L.B., Peet D.J., Fujii N., Gronthos S., Harris A.L., Zannettino A.C. Hypoxia-inducible factor-2 is a novel regulator of aberrant CXCL12 expression in multiple myeloma plasma cells.
Haematologica **95**(5) 776-784. (2010)
30. Mandawat A., Fiskus W., Buckley K.M., Robbins K., Rao R., Balusu R., Navenot J.M., Wang Z.X., Ustun C., Chong D.G., Atadja P., Fujii N., Peiper S.C., Bhalla K. Pan-histone deacetylase (HDAC) inhibitor panobinostat depletes CXCR4 levels and signaling and exerts synergistic anti-myeloid activity in

- combination with CXCR4 antagonists.
Blood **116**(24) 5306-5315. (2010)
31. Omatsu Y., Sugiyama T., Kohara H., Kondoh G., Fujii N., Kohno K., Nagasawa T.
The essential functions of adipo-osteogenic progenitors as the hematopoietic stem and progenitor cell niche.
Immunity **33**(3) 387-399. (2010)
32. Oishi S., Watanabe K., Ito S.; Tanaka M., Nishikawa H., Ohno H., Shimane K., Izumi K., Sakagami Y., Kodama E. N., Matsuoka M., Asai A., Fujii N.
Affinity selection and sequence-activity relationships of HIV-1 membrane fusion inhibitors directed at the drug-resistant variants.
Med. Chem. Commn. in press.
33. Ohta Y., Tokimizu Y., Oishi S., Fujii N., Ohno H.
Direct Synthesis of Quinazolines through Copper-catalyzed Reaction of Aniline-Derived Benzamidines.
Org. Lett. **12**(17) 3963-3965 (2010)
34. Ding S., Nishizawa K., Kobayashi T., Oishi S., Lv J., Fujii N., Ogawa O., Nishiyama H.
A Potent Chemotherapeutic Strategy for Bladder Cancer: (S)-Methoxy-Trityl-L-Cysteine, a Novel Eg5 Inhibitor.
J. Urol. **184**(3) 1175-1181 (2010)
35. Oishi S., Watanabe T., Sawada J., Asai A., Ohno H., Fujii N.
Kinesin spindle protein (KSP) inhibitors with 2,3-fused indole scaffolds.
J. Med. Chem. **53**(13) 5054-5058. (2010)
36. Inuki S., Yoshimitsu Y., Oishi S., Fujii N., Ohno H.
Ring-Construction/Stereoselective Functionalization Cascade: Total Synthesis of Pachastrissamine (Jaspine B) through Palladium-Catalyzed Bis-cyclization of Propargyl Chlorides and Carbonates.
J. Org. Chem. **75**(11) 3831-3842. (2010)
37. Izumi K., Nakamura S., Nakano H., Shimura K., Sakagami Y., Oishi S., Uchiyama S., Ohkubo T., Kobayashi Y., Fujii N., Matsuoka M., Kodama E.N.
Characterization of HIV-1 resistance to a fusion inhibitor, N36, derived from the gp41 amino terminal heptad repeat.
Antiviral Res. **87**(2) 179186 (2010)
38. Yoshimitsu Y., Inuki S., Oishi S., Fujii N., Ohno H.
Stereoselective divergent synthesis of four diastereomers of pachastrissamine

- (jaspine B).
- J. Org. Chem.* **75** 3843-3846. (2010)
39. Okano A., Oishi S., Tanaka T., Fujii N., Ohno H.
Construction of linked nitrogen heterocycles by palladium(0)-catalyzed intramolecular domino cyclization of 2-alkynylaziridines bearing a 2-aminoethyl group via ring expansion with isocyanate.
J. Org. Chem. **75** 3396-3400. (2010)
40. Okano A., Tsukamoto K., Kosaka S., Maeda H., Oishi S., Tanaka T., Fujii N., Ohno H.
Synthesis of fused and linked bicyclic nitrogen heterocycles by palladium-catalyzed domino cyclization of propargyl bromides.
Chem. Eur. J. in press (2010)
41. Nishizawa K., Nishiyama H., Ohishi S., Tanahara N., Kotani H., Mikami Y., Toda Y., Evans BJ., Peiper SC., Saito R., Watanabe J., Fujii N., Ogawa O.
Fluorescent imaging of high grade bladder cancer using a specific antagonist for chemokine receptor CXCR4.
Int J Cancer. **127**(5) 1180-1187. (2010)
42. Melchionna R., Di Carlo A., De Mori R., Cappuzzello C., Barberi L., Musaro A., Cencioni C., Fujii N., Tamamura H., Crescenzi M., Capogrossi MC., Napolitano M., Germani A.
Induction of myogenic differentiation by SDF-1 via CXCR4 and CXCR7 receptors.
Muscle Nerve. **41** 828-835. (2010)
43. Hirano K., Inaba Y., Watanabe T., Oishi S., Fujii N., Ohno H.
Gold-catalyzed intramolecular alkyne cycloisomerization cascade: direct synthesis of aryl-annulated[a]carbazoles from aniline-substituted diethynylarenes.
Adv. Syn. Catal. **352**(2) 368-372. (2010)
44. Narumi T., Hayashi R., Tomita K., Kobayashi K., Tanahara N., Ohno H., Naito T., Kodama E., Matsuoka M., Oishi S., Fujii N.
Synthesis and biological evaluation of selective CXCR4 antagonists containing alkene dipeptide isosteres.
Org. Biomol. Chem. **8**(3) 616-621 (2010)
45. Mizuhara T., Oishi S., Fujii N., Ohno H.
Efficient synthesis of pyrimido[1,2-c][1,3]benzothiazin-6-imines and related tricyclic heterocycles by SNAr-Type C-S, C-N, or C-O bond formation with

heterocumulenes.

J. Org. Chem. **75**(1) 265-268 (2010)

46. Kajiwara K., Watanabe K., Tokiwa R., Kurose T., Ohno H., Tsutsumi H., Hata Y., Izumi K., Kodama E., Matsuoka M., Oishi S., Fujii N.
Bioorganic synthesis of a recombinant HIV-1 fusion inhibitor, SC35EK, with an N-terminal pyroglutamate capping group.
Bioorg. Med. Chem. **17**(23) 7964-7970 (2009)
47. Tanaka M., Kajiwara K., Tokiwa R., Watanabe K., Ohno H., Tsutsumi H., Hata Y., Izumi K., Kodama E., Matsuoka M., Oishi S., Fujii N.
Bioorganic synthesis of end-capped anti-HIV peptides by simultaneous cyanocysteine-mediated cleavages of recombinant proteins.
Bioorg. Med. Chem. **17**(21) 7487-7492 (2009)
48. Ohta Y., Chiba H., Oishi S., Fujii N., Ohno H.
Construction of nitrogen heterocycles bearing an aminomethyl group by copper-catalyzed domino three-component coupling-cyclization.
J. Org. Chem. **74**(18) 7052-7058 (2009)
49. Inuki S., Yoshimitsu Y., Oishi S., Fujii N., Ohno H.
Ring-Construction/Stereoselective Functionalization Cascade: Total Synthesis of Pachastrissamine (Jaspine B) through Palladium-Catalyzed Bis-cyclization of Bromoallenes.
Org. Lett. **11**(19) 4478-4481 (2009)
50. Watabe T., Terakawa Y., Watanabe K., Ohno H., Nakano H., Nakatsu T., Kato H., Izumi K., Kodama E., Matsuoka M., Kitaura K., Oishi S., Fujii N.
X-ray crystallographic study of an HIV-1fusion inhibitor with the gp41 S138A substitution.
J. Mol. Biol. **392**(3) 657-665 (2009)
51. Arthur A., Shi S., Zannettino A. C., Fujii N., Gronthos S., Koblar S. A.
Implanted adult human dental pulp stem cells induce endogenous axon guidance.
Stem Cells. **27**(9) 2229-2237 (2009)
52. Ohta Y., Kubota Y., Watabe T., Chiba H., Oishi S., Fujii N., Ohno H.
Rapid access to 3-(aminomethyl)isoquinoline-fused polycyclic compounds by copper-catalyzed four-component coupling, cascade cyclization, and oxidation.
J. Org. Chem. **74**(16) 6299-6302 (2009)
53. Tanaka T., Nomura W., Narumi T., Esaka A., Oishi S., Ohashi N., Itotani K., Evans B. J., Wang Z., Peiper S. C., Fujii N., Tamamura H.

Structure-activity relationship study on artificial CXCR4 ligands possessing the cyclic pentapeptide scaffold: the exploration of amino acid residues of pentapeptides by substitutions of several aromatic amino acids.

Org. Biomol. Chem. **7**(18) 3805-3809 (2009)

54. Oishi S., Kodera Y., Nishikawa H., Kamitani H., Watabe T., Ohno H., Tochikura T., Shimane K., Kodama E., Matsuoka M., Mizukoshi F., Tsujimoto H., Fujii N. Design and synthesis of membrane fusion inhibitors against the feline immunodeficiency virus.
Bioorg. Med. Chem. **17**(14) 4916-4920 (2009)
55. Evans B.J., Wang Z., Broach J.R., Oishi S., Fujii N., Peiper S.C. Expression of CXCR4, a G-protein-coupled receptor for CXCL12 in yeast identification of new-generation inverse agonists.
Methods Enzymol. **460** 399-412 (2009)
56. Ueno M., Kodama E. N., Shimura K., Sakurai Y., Kajiwara K., Sakagami Y., Oishi S., Fujii N., Matsuoka M..
Synonymous mutations in stem-loop III of Rev responsive elements enhance HIV-1 replication impaired by primary mutations for resistance to enfuvirtide.
Antiviral Res. **82**(1) 67-72 (2009)
57. Ohta Y., Oishi S., Fujii N., Ohno H.
Facile synthesis of 1,2,3,4-tetrahydro-β-carbolines by one-pot domino three-component indole formation and nucleophilic cyclization.
Org. Lett. **11**(9) 1979-1982 (2009)
58. Watanabe T., Oishi S., Fujii N., Ohno H.
Palladium-catalyzed direct synthesis of carbazoles via one-pot N-arylation and oxidative biaryl coupling: synthesis and mechanistic study.
J. Org. Chem. **74**(13) 4270-4276 (2009)
59. Suzuki Y., Ohta Y., Oishi S., Fujii N., Ohno H.
Efficient synthesis of aminomethylated pyrroloindoles and dipyrrolopyridines via controlled copper-catalyzed domino multi-component coupling and bis-cyclization.
J. Org. Chem. **74**(11) 4246-4251 (2009)
60. Kobayashi K., Narumi T., Oishi S., Ohno H., Fujii N.
Amino acid-based synthesis of trifluoromethylalkene dipeptide isosteres by alcohol-assisted nucleophilic trifluoromethylation and organozinc-copper-mediated SN2' alkylation.
J. Org. Chem. **74**(12) 4626-4629 (2009)

61. Oishi S., Kamitani H., Kodera Y., Watanabe K., Kobayashi K., Narumi T., Tomita K., Ohno H., Naito T., Kodama E., Matsuoka M., Fujii N.
Peptide bond mimicry by (*E*)-alkene and (Z)-fluoroalkene peptide isosteres: synthesis and bioevaluation of α -helical anti-HIV peptide analogues.
Org. Biomol. Chem. **7**(14) 2872-2877 (2009)
62. Mizuhara T., Inuki S., Oishi S., Fujii N., Ohno H.
 Cu(II) -mediated oxidative intermolecular ortho C-H functionalisation using tetrahydropyrimidine as the directing group.
Chem. Commun. (23) 3413-3415 (2009)
63. Navenot J.M., Fujii N., Peiper S.C.
KiSS1 Metastasis Suppressor Gene Product Induces Suppression of Tyrosine Kinase Receptor Signaling to Akt, TNF Family Ligand Expression, and Apoptosis.
Mol. Pharmacol. **75**(5) 1074-1083 (2009)
64. Navenot J.M., Fujii N., Peiper S.C.
Activation of Rho and ROCK by GPR54 and KiSS1 Metastasis Suppressor Gene Product Induces Changes of Cell Morphology and Contributes to Apoptosis.
Mol. Pharmacol. **75**(6) 1300-1306 (2009)
65. Diamond P., Labrinidis A., Martin S.K., Farrugia A.N., Gronthos S., To L.B., Fujii N., O' Loughlin P.D., Evdokiou A., Zannettino A.C.W.
Targeted disruption of the CXCL12/CXCR4 axis inhibits osteolysis in a murine model of myeloma-associated bone loss.
J. Bone Miner. Res. **24**(7) 1150-1161 (2009)
66. Mizukoshi F., Baba K., Goto-Koshino Y., Setoguchi-Mukai A., Fujino Y., Ohno K., Tamamura H., Oishi S., Fujii N., Tsujimoto H.
Inhibitory effect of newly developed CXC-chemokine receptor 4 antagonists on the infection with feline immunodeficiency virus.
Vet. Microbiol. **71**(1) 121-124 (2009)
67. Naito T., Izumi K., Kodama E., Sakagami Y., Kajiwara K., Nishikawa H., Watanabe K., Sarafianos S.G., Oishi S., Fujii N., Matsuoka M.
SC29EK, a peptide fusion inhibitor with enhanced α - helicity, inhibits replication of human immunodeficiency virus type 1 mutants resistant to enfuvirtide.
Antimicrob. Agents Chemother. **53**(3) 1013-1018 (2009)
68. Izumi K., Kodama E., Shimura K., Sakagami Y., Watanabe K., Ito S., Watabe T., Terakawa Y., Nishikawa H., Sarafianos S. G., Kitaura K., Oishi S., Fujii N., Matsuoka M.

- Design of peptide- based inhibitors of HIV-1 strains resistant to T-20.
J. Biol. Chem. **284**(8) 4914-4920 (2009)
69. Mizukoshi F., Baba F., Goto Y., Setoguchi A., Fujino Y., Ohno K., Oishi S., Kodera Y., Fujii N., Tsujimoto H.
Antiviral activity of membrane fusion inhibitors that target gp40 of the feline immunodeficiency virus envelope protein.
J. Vet. Med. Sci. **71**(1) 121-124 (2009)
70. Kitaori T., Ito H., Schwarz E.M., Tsutsumi R., Yoshitomi H., Oishi S., Nakano M., Fujii N., Nagasawa T., Nakamura T.
SDF-1/CXCR4 signaling is critical for the recruitment of mesenchymal stem cells to the fracture site during skeletal repair.
Arthritis Rheum. **60**(3) 813-823 (2009)
71. Nishikawa H., Nakamura S., Kodama E., Ito S., Kajiwara K., Izumi K., Sakagami Y., Oishi S., Ohkubo T., Kobayashi Y., Otaka A., Fujii N., Matsuoka M.
Electrostatically constrained α -helical peptide inhibitors replication of HIV-1 resistant to enfuvirtide.
Int. J. Biochem. Cell Biol. **41**(4) 891-899 (2009)
72. Yamaki Y., Shigenaga A., Tomita K., Narumi T., Fujii N., Otaka A.
Synthesis of fluoroalkene dipeptide isosteres by intramolecular redox reaction utilizing N-heterocyclic carbenes (NHCs).
J. Org. Chem. **74** 3272-3277 (2009)
73. Katagiri F., Nagai K., Kida A., Tomita K., Oishi S., Takeyama M., Doi R., Fujii N.
Clinical significance of plasma metastin level in pancreatic cancer patients.
Oncol. Rep. **21**(3) 815-819 (2009)
74. Tomita K., Popiel H.A., Nagai Y., Toda T., Yoshimitsu Y., Ohno H., Oishi S., Fujii N
Structure-activity relationship study on polyglutamine binding peptide QBP1.
Bioorg. Med. Chem. **17**(3) 1259-1263 (2009)
75. Nagai K., Doi R., Katagiri F., Ito T., Kida A., Koizumi M., Masui T., Kawaguchi Y., Tomita K., Oishi S., Fujii N., Uemoto S.
Prognostic value of metastin expression in human pancreatic cancer.
J. Exp. Clin. Cancer Res. **28**(9) (2009)
76. Evans B.J., Wang Z., Mobley L., Khosravi D., Fujii N., Navenot J.M., Peiper S.C.
Physical association of GPR54 C-terminal with protein phosphatase 2A.
Biochem. Biophys. Res. Commun. **377**(4) 1067-1071 (2008)

77. Tanaka T., Tsutsumi H., Nomura W., Tanabe Y., Ohashi N., Esaka A., Ochiai C., Sato J., Itotani K., Murakami T., Ohba K., Yamamoto N., Fujii N., Tamamura H. Structure-activity relationship study of CXCR4 antagonists bearing the cyclic pentapeptide scaffold: identification of the new pharmacophore.
Org. Biomol. Chem. **6**(23) 4374-4377 (2008)
78. Tomita K., Oishi S., Ohno H., Peiper S.C., Fujii N. Development of novel GPR54 agonists with resistance to degradation by matrix metalloproteinase.
J. Med. Chem. **51**(23) 7645-7649 (2008)
79. Nishikawa H., Oishi S., Fujita M., Watanabe K., Tokiwa R., Ohno H., Kodama E., Izumi K., Kajiwara K., Naitoh T., Matsuoka M., Otaka A., Fujii N. Identification of minimal sequence for HIV-1 fusion inhibitors.
Bioorg. Med. Chem. **16**(20) 9184-9187 (2008)
80. Inuki S., Oishi S., Fujii N., Ohno H. Total synthesis of (\pm)-lysergic acid, lysergol, and isolysergol by palladium-catalyzed domino cyclization of amino allenes bearing a bromoindolyl group.
Org. Lett. **10**(22) 5239-5242 (2008)
81. Mikami S., Nakase H., Yamamoto S., Takeda Y., Yoshino T., Kasahara K., Ueno S., Uza N., Oishi S., Fujii N., Nagasawa T., Chiba T. Blockade of CXCL12/CXCR4 axis ameliorates murine experimental colitis.
J. Pharmacol. Exp. Ther. **327**(2) 383-392 (2008)
82. Ohta Y., Chiba H., Oishi S., Fujii N., Ohno H. Concise synthesis of indole-fused 1,4-diazepines through copper(I)-catalyzed domino three-component coupling-cyclization-N-arylation under microwave irradiation.
Org. Lett. **10**(16) 3535-3538 (2008)
83. Yano Y., Yano A., Oishi S., Sugimoto Y., Tsujimoto G., Fujii N., Matsuzaki K. Coiled-coil tag-probe system for quick labeling of membrane receptors in living cells.
ACS Chem. Biol. **3**(6) 341-345 (2008)
84. Ueda S., Kato M., Inuki S., Ohno H., Evans B., Wang Z., Peiper S.C., Izumi K., Kodama E., Matsuoka M., Nagasawa H., Oishi S., Fujii N. Identification of novel nonpeptide CXCR4 antagonists by ligand-based design approach.
Bioorg. Med. Chem. Lett. **18**(14) 4124-4129 (2008)

85. Okano A., Mizutani T., Oishi S., Tanaka T., Ohno H., Fujii N.
Palladium-catalysed biscyclisation of allenic bromoalkenes through zipper-mode cascade.
Chem. Commun. **30** 3534-3536 (2008)
86. Nishikawa H., Kodama E., Sakakibara A., Fukudome A., Izumi K., Oishi S., Fujii N., Matsuoka M.
Novel screening systems for HIV-1 fusion mediated by two extra-virion heptad repeats of gp4.
Antivir. Res. **80**(1) 71-76(2008)
87. Nakata H., Steinberg S.M., Koh Y., Maeda K., Takaoka Y., Tamamura H., Fujii N., Mitsuya H.
Potent synergistic anti-HIV effects using combinations of a CCR5.
Antimicrob. Agents. Chemother. **52**(6) 2111-2119 (2008)
88. Watanabe T., Oishi S., Fujii N., Ohno H.
Palladium-catalyzed sp³ C – H activation of simple alkyl groups: direct preparation of indoline derivatives from N-alkyl-2-bromoanilines.
Org. Lett. **10**(9) 1759-1762 (2008)
89. Inokuchi E., Narumi T., Niida A., Kobayashi K., Tomita K., Oishi S., Ohno H., Fujii N.
Efficient synthesis of trifluoromethyl and related trisubstituted alkene dipeptide isosteres by palladium-catalyzed carbonylation of amino acid-derived allylic carbonates.
J. Org. Chem. **73**(10) 3942-3945 (2008)
90. Narumi T., Tomita K., Inokuchi E., Kobayashi K., Oishi S., Ohno H., Fujii N.
Diastereoselective synthesis of highly functionalized fluoroalkene dipeptide isosteres and its application to Fmoc-based solid phase synthesis of a cyclic pentapeptide mimetic.
Tetrahedron, **64**(19) 4332-4346 (2008)
91. Suzuki Y., Cluzeau J., Hara T., Hirasawa A., Tsujimoto G., Oishi S., Ohno H., Fujii N.
Structure-activity relationship of pyrazine-based CK2 inhibitors: synthesis and biological evaluation of 2,6-disubstituted pyrazine and 4,6-disubstituted pyrimidine derivatives.
Archiv. Pharm. (Weinheim) **341**(9) 554-561 (2008)
92. Ohno H., Okano A., Kosaka S., Tsukamoto K., Ohata M., Ishihara K., Maeda H., Tanaka T., Fujii N.

- Direct construction of bicyclic heterocycles by palladium-catalyzed domino cyclization of propargyl bromides.
Org. Lett. **10**(6) 1171-1174 (2008)
93. Tomita K., Oishi S., Ohno H., Fujii N.
Structure-activity relationship study and NMR analysis of fluorobenzoyl pentapeptide GPR54 agonists.
Biopolymers, **90**(4) 503-511 (2008)
94. Driessens W.H., Fujii N., Tamamura H., Sullivan S.M.
Development of peptide-targeted lipoplexes to CXCR4-expressing rat glioma cells and rat proliferating endothelial cells.
Mol. Ther. **16**(3) 516-524 (2008)
95. Ohta Y., Oishi S., Fujii N., Ohno H.
Facile synthesis of 3-(aminomethyl)isoquinolines by copper-catalysed domino four-component coupling and cyclisation.
Chem. Commun. **7** 835-837 (2008)
96. Oishi S., Masuda R., Evans B., Ueda S., Goto Y., Ohno H., Hirasawa A., Tsujimoto G., Wang Z., Peiper S.C., Naito T., Kodama E., Matsuoka M., Fujii N.
Synthesis and application of fluorescein- and biotin-labeled molecular probes for chemokine receptor CXCR4.
ChemBioChem, **9**(7) 1154-1158 (2008)
97. Kasiyanov A., Fujii N., Tamamura H., Xiong H.
Modulation of network-driven, GABA-mediated giant depolarizing potentials by SDF-1 α in the developing hippocampus.
Dev. Neurosci. **30**(4) 285-292 (2008)
98. Oishi S., Ito S., Nishikawa H., Watanabe K., Tanaka M., Ohno H., Izumi K., Sakagami Y., Kodama E., Matsuoka M., Fujii N.
Design of a novel HIV-1 fusion inhibitor that displays a minimal interface for binding affinity.
J. Med. Chem. **51**(3) 388-391 (2008)
99. Liapi A., Pritchett J., Jones O., Fujii N., Parnavelas J.G., Nadarajah B.
Stromal-derived factor 1 signalling regulates radial and tangential migration in the developing cerebral cortex.
Dev. Neurosci. **30**(1-3) 117-131 (2008)
100. Ujike M., Nishikawa H., Otaka A., Yamamoto N., Yamamoto N., Matsuoka M., Kodama E., Fujii N., Taguchi F.
Heptad repeat-derived peptides block the protease-mediated direct entry from

- cell surface of SARS coronavirus but not entry via endosomal pathway.
J. Virol. **82**(1) 588-592 (2008)
101. Sasaki Y., Fujii N., Otaka A.
Development of copper-mediated allylation of γ -activated- α , β -unsaturated lactam toward peptide mimetic synthesis.
Tetrahedron lett. **48**(18) 3221-3224 (2007)
102. Watanabe T., Ueda S., Inuki S., Oishi S., Fujii N., Ohno H.
One-pot synthesis of carbazoles by palladium-catalyzed *N*-arylation and oxidative coupling.
Chem. Commun. (43) 4516-4518 (2007)
103. Watanabe T., Oishi S., Fujii N., Ohno H.
Gold-catalyzed hydroarylation of allenes: a highly regioselective carbon-carbon bond formation producing six-membered rings.
Org. Lett. **9**(23) 4821-4824 (2007)
104. Ohno H., Iuchi M., Fujii N., Tanaka T.
Zipper-mode double C-H activation: palladium-catalyzed direct construction of highly-fused heterocyclic systems.
Org. Lett. **9**(23) 4813-4815(2007)
105. Kohara H., Omatsu Y., Sugiyama T., Noda M., Fujii N., Nagasawa T.
Development of plasmacytoid dendritic cells in bone marrow stromal cell niches requires CXCL12-CXCR4 chemokine signaling.
Blood, **110**(13) 4153-4160(2007)
106. Kabashima K., Shiraishi N., Sugita K., Mori T., Onoue A., Kobayashi M., Sakabe JI., Yoshiki R., Tamamura H., Fujii N., Inaba K., Tokura Y.
CXCL12-CXCR4 engagement is required for migration of cutaneous dendritic cells.
Am. J. Pathol. **171**(4) 1249-1257(2007)
107. Zhang W., Navenot J.M., Frilot N.M., Fujii N., Peiper S.C.
Association of nucleophosmin negatively regulates CXCR4-mediated G protein activation and chemotaxis.
Mol. Pharmacol. **72**(5) 1310-1321 (2007)]
108. Tamamura H., Tanaka T., Tsutsumi H., Nemoto K., Mizokami S., Ohashi N., Oishi S., Fujii N.
Versatile use of acid-catalyzed ring-opening ofβ-aziridinyl-α,β-enoates to stereoselective synthesis of peptidomimetics.
Tetrahedron, **63**(37) 9243-9254 (2007)

109. Kabashima K., Sugita K., Shiraishi N., Tamamura H., Fujii N., Tokura Y.
CXCR4 engagement promotes dendritic cell survival and maturation.
Biochem. Biophys. Res. Commun. **361**(4) 1012-1016 (2007)
110. Narumi T., Tomita K., Inokuchi E., Kobayashi K., Oishi S., Ohno H., Fujii N.
Facile synthesis of fluoroalkenes by palladium-catalyzed reductive defluorination of allylic gem-difluorides.
Org. Lett. **9**(17) 3465-3468 (2007)
111. Ohno H., Aso A., Kadoh Y., Fujii N., Tanaka T.
Heck-type cyclization of oxime ethers: stereoselective carbon-carbon bond formation with aryl halides to produce heterocyclic oximes.
Angew. Chem. Int. Ed. **46**(33) 6325-6328 (2007)
112. Kubonishi S., Kikuchi T., Yamaguchi S., Tamamura H., Fujii N., Watanabe T., Arenzana-Seisdedos F., Ikeda K., Matsui T., Tanimoto M., Katayama Y.
Rapid hematopoietic progenitor mobilization by sulfated colominic acid.
Biochem. Biophys. Res. Commun. **355**(4) 970-975 (2007)
113. Cluzeau J., Oishi S., Ohno H., Wang Z., Evans B., Peiper S.C., Fujii N.
Design and synthesis of all diastereomers of cyclic pseudo-dipeptides as mimics of cyclic CXCR4 pentapeptide antagonists.
Org. Biomol. Chem. **5**(12) 1915-1923 (2007)
114. Ohno H., Mizutani T., Kadoh Y., Aso A., Miyamura K., Fujii N., Tanaka T.
A highly regio- and stereoselective formation of bicyclo[4.2.0]oct-5-ene derivatives through thermal intramolecular [2 + 2] cycloaddition of allenes.
J. Org. Chem. **72**(12) 4378-4389 (2007)
115. Ono Y., Kashiwagi H., Esaki T., Tadakatsu T., Sato H., Fujii N.
Systematic solution-phase parallel synthesis of active vitamin D₃ analogs with elongated side chains and their cell differentiation activities.
J. Comb. Chem. **9**(4) 711-716 (2007)
116. Ogo N., Oishi S., Matsuno K., Sawada J., Fujii N., Asai A.
Synthesis and biological evaluation of L-cysteine derivatives as mitotic kinesin Eg5 inhibitors.
Bioorg. Med. Chem. Lett. **17**(14) 3921-3924 (2007)
117. Tomita K., Oishi S., Cluzeau J., Ohno H., Navenot J.M., Wang Z.X., Peiper S.C., Akamatsu M., Fujii N.
SAR and QSAR studies on the N-terminally acylated pentapeptide agonists for GPR54.
J. Med. Chem. **50**(14) 3222-3228 (2007)

118. Katagiri F., Tomita K., Oishi S., Takeyama M., Fujii N.
Establishment and clinical application of enzyme immunoassays for determination of luteinizing hormone releasing hormone and metastin.
J. Pept. Sci. **13**(6) 422-429 (2007)
119. Juarez J., Dela Pena A., Baraz R., Hewson J., Khoo M., Cisterne A., Fricker S., Fujii N., Bradstock K.F., Bendall L.J.
CXCR4 antagonists mobilize childhood acute lymphoblastic leukemia cells into the peripheral blood and inhibit engraftment.
Leukemia, **21**(6) 1249-1257 (2007)
120. Berchiche Y.A., Chow K.Y., Lagane B., Leduc M., Percherancier Y., Fujii N., Tamamura H., Bachelerie F., Heveker N.
Direct assessment of CXCR4 mutant conformations reveals complex link between receptor structure and G α i activation.
J. Biol. Chem. **282**(8) 5111-5115 (2007)
121. Tomita K., Narumi T., Niida A., Oishi S., Ohno H., Fujii N.
Fmoc-based solid-phase synthesis of GPR54-agonistic pentapeptide derivatives containing alkene- and fluoroalkene-dipeptide isosteres.
Biopolymers, **88**(2) 272-278 (2007)
122. Ohno H., Ohta Y., Oishi S., Fujii N.
Direct synthesis of 2-(aminomethyl)indoles through copper(I)-catalyzed domino three-component coupling and cyclization reactions.
Angew. Chem. Int. Ed. **46**(13) 2295-2298 (2007)
123. Ueda S., Oishi S., Wang Z.X., Araki T., Tamamura H., Cluzeau J., Ohno H., Kusano S., Nakashima H., Trent J.O., Peiper S.C., Fujii N.
Structure-activity relationships of cyclic peptide-based chemokine receptor CXCR4 antagonists: disclosing the importance of side-chain and backbone functionalities.
J. Med. Chem. **50**(2) 192-198 (2007)
124. Hamaguchi H., Kosaka S., Ohno H., Fujii N., Tanaka T.
Bromoallenes as allyl dication equivalents in the presence or absence of palladium(0): direct construction of bicyclic sulfamides containing five- to eight-membered rings by tandem cyclization of bromoallenes.
Chem. Eur. J. **13**(6) 1692-1708 (2007)
125. Tanaka M., Oishi S., Ohno H., Fujii N.
A novel oxazolidine linker for the synthesis of peptide aldehydes.
Int. J. Pep. Res. Ther. **13**(1-2) 271-279(2007)

126. Sasaki Y., Shigenaga A., Fujii N., Otaka A.
Synthesis of (\mathcal{Z})-alkene-containing *cis*-proline dipeptide mimetics using samarium (II) diiodide (SmI_2)-mediated reductive-alkylation reaction.
Tetrahedron, **63**(9) 2000-2008(2007)
127. Kang S.-U., Choi W. J., Oishi S., Lee K., Karki R.G., Worthy K.M., Bindu L.K., Nicklaus M.C., Fisher R.J., Burke T.R.Jr.
Examination of acylated 4-aminopiperidine-4-carboxylic acid residues in the pTyr+1 position of Grb2 SH2 domain-binding tripeptides.
J. Med. Chem. **50**(8) 1978-1982(2007)
128. Narumi T., Niida A., Tomita K., Oishi S., Otaka A., Ohno H., Fujii N.
A novel one-pot reaction involving organocopper-mediated reduction/transmetalation/asymmetric alkylation, leading to the diastereoselective synthesis of functionalized (\mathcal{Z})-fluoroalkene dipeptide isosteres.
Chem. Commun. (45) 4720-4722 (2006)
129. Tomita K., Niida A., Oishi S., Ohno H., Cluzeau J., Navenot J.M., Wang Z.X., Peiper S.C., Fujii N.
Structure-activity relationship study on small peptidic GPR54 agonists.
Bioorg. Med. Chem. **14**(22) 7595-7603 (2006)
130. Sasaki Y., Niida A., Tsuji T., Shigenaga A., Fujii N., Otaka A.
Stereoselective synthesis of (\mathcal{Z})-alkene-containing proline dipeptide mimetics.
J. Org. Chem. **71**(13) 4969-4979 (2006)
131. Niida A., Tanigaki H., Inokuchi E., Sasaki Y., Oishi S., Ohno H., Tamamura H., Wang Z., Peiper S.C., Kitaura K., Otaka A., Fujii N.
Stereoselective synthesis of 3,6-disubstituted-3,6-dihydropyridin-2-ones as potential diketopiperazine mimetics using organocopper-mediated anti-Sn_{2'} reactions and their use in the preparation of low-molecule CXCR4 antagonists.
J. Org. Chem. **71**(10) 3942-3951 (2006)
132. Niida A., Mizumoto M., Narumi T., Inokuchi E., Oishi S., Ohno H., Otaka A., Kitaura K., Fujii N.
Synthesis of (\mathcal{Z})-alkene and (E)-fluoroalkene-containing diketopiperazine mimetics utilizing organocopper-mediated reduction-alkylation and diastereoselectivity examination using DFT calculations.
J. Org. Chem. **71**(11) 4118-4129 (2006)
133. Hanaoka H., Mukai T., Tamamura H., Mori T., Ishino S., Ogawa K., Iida Y., Doi R., Fujii N., Saji H.

- Development of a ^{111}In -labeled peptide derivative targeting a chemokine receptor, CXCR4, for imaging tumors.
Nucl. Med. Biol. **33**(4) 489-494 (2006)
134. Tamamura H., Ojida A., Ogawa T., Tsutsumi H., Masuno H., Nakashima H., Yamamoto N., Hamachi I., Fujii N.
Identification of a new class of low molecular weight antagonists against the chemokine receptor CXCR4 having the dipicolylamine-zinc(II) complex structure.
J. Med. Chem. **49**(11) 3412-3415 (2006)
135. Tamamura H., Tsutsumi H., Masuno H., Mizokami S., Hiramatsu K., Wang Z., Trent J.O., Nakashima H., Yamamoto N., Peiper S.C., Fujii N.
Development of a linear type of low molecular weight CXCR4 antagonists based on T140 analogs.
Org. Biomol. Chem. **4**(12) 2354-2357 (2006)
136. Tsuda M., Terada T., Irie M., Katsura T., Niida A., Tomita K., Fujii N., Inui K.
Transport characteristics of a novel PEPT1 substrate, antihypotensive drug midodrine, and its amino acid derivatives.
J. Pharmacol. Exp. Ther. **318**(1) 455-460 (2006)
137. Menu E., Asosingh K., Indraccolo S., De Raeve H., Van Riet I., Van Valckenborgh E., Van de Broek I., Fujii N., Tamamura H., Van Camp B., Vanderkerken K.
The involvement of stromal derived factor 1 α in homing and progression of multiple myeloma in the 5TMM model.
Haematologica, **91**(5) 605-612 (2006)
138. Kasyanov A., Tamamura H., Fujii N., Xiong H.
HIV-1 gp120 enhances giant depolarizing potentials via chemokine receptor CXCR4 in neonatal rat hippocampus.
Eur. J. Neurosci. **23**(5) 1120-1128 (2006)
139. Niida A., Tomita K., Mizumoto M., Tanigaki H., Terada T., Oishi S., Otaka A., Inui K., Fujii N.
Unequivocal synthesis of (\mathcal{D})-alkene and (\mathcal{E})-fluoroalkene dipeptide isosteres to probe structural requirements of the peptide transporter PEPT1.
Org. Lett. **8**(4) 613-616 (2006)
140. Ohta Y., Itoh S., Shigenaga A., Shintaku S., Fujii N., Otaka A.
Cysteine-derived *S*-protected oxazolidinones: potential chemical devices for the preparation of peptide thioesters.

- Org. Lett.* **8**(3) 467-470 (2006)
141. Ohno H., Kadoh Y., Fujii N., Tanaka T.
Potassium carbonate-promoted stereospecific 5-endo-trig cyclization of unactivated allenes in the absence of any transition metals.
- Org. Lett.* **8**(5) 947-950 (2006)
142. Oishi S., Miyamoto K., Niida A., Yamamoto M., Ajito K., Tamamura H., Otaka A., Kuroda Y., Asai A., Fujii N.
Application of tri- and tetrasubstituted alkene dipeptide mimetics to conformational studies of cyclic RGD peptides.
- Tetrahedron*, **62**(7) 1416-1424 (2006)
143. Avniel S., Arik Z., Maly A., Sagie A., Basst H.B., Yahana M.D., Weiss I.D., Pal B., Wald O., Ad-El D., Fujii N., Arenzana-Seisdedos F., Jung S., Galun E., Gur E., Peled A.
Involvement of the CXCL12/CXCR4 pathway in the recovery of skin following burns.
- J. Invest. Dermatol.* **126**(2) 468-476 (2006)
144. Dewan M.Z., Uchihara J.N., Terashima K., Honda M., Sata T., Ito M., Fujii N., Uozumi K., Tsukasaki K., Tomonaga M., Kubuki Y., Okayama A., Toi M., Mori N., Yamamoto N.
Efficient intervention of growth and infiltration of primary adult T-cell leukemia cells by an HIV protease inhibitor, ritonavir.
- Blood*, **107**(2) 716-724 (2006)
145. Niida A., Wang Z., Tomita K., Oishi S., Tamamura H., Otaka A., Navenot J.M., Broach J.R., Peiper S.C., Fujii N.
Design and synthesis of downsized metastin (45-54) analogs with maintenance of high GPR54 agonistic activity.
- Bioorg. Med. Chem. Lett.* **16**(1) 134-137 (2006)
146. Navenot J.M., Wang Z., Chopin M., Fujii N., Peiper S.C.
Kisspeptin-10-induced signaling of GPR54 negatively regulates chemotactic responses mediated by CXCR4: a potential mechanism for the metastasis suppressor activity of kisspeptins.
- Cancer Res.* **65**, 10450-10456(2005)
147. Tamamura H., Esaka A., Ogawa T., Araki T., Ueda S., Wang Z., Trent J.O., Tsutsumi H., Masuno H., Nakashima H., Yamamoto N., Peiper S.C., Otaka A., Fujii N.
Structure-activity relationship studies on CXCR4 antagonists having cyclic

- pentapeptide scaffolds.
Org. Biomol. Chem. **3**, 4392-4394 (2005)
148. Zhong Y., Yoshinaka Y., Takeda T., Shimizu N., Yoshizaki S., Inagaki Y., Matsuda S., Honda G., Fujii N., Yamamoto N.
Highly potent anti-HIV-1 activity isolated from fermented Polygonum tinctorium Aiton.
Antivir. Res. **66**(2-3) 119-128 (2005)
149. Retz M., Sidhu S.S., Lehmann J., Tamamura H., Fujii N., Basbaum C.
New HIV-drug inhibits in vitro bladder cancer migration and invasion.
Eur. Urol. **48**(6) 1025-1030 (2005)
150. Ueda S., Fujita M., Tamamura H., Fujii N., Otaka A.
Photolabile protection for one-pot sequential native chemical ligation.
Chembiochem, **6**(11) 1983-1986 (2005)
151. Hartmann TN., Burger J.A., Glodek A., Fujii N., Burger M.
CXCR4 chemokine receptor and integrin signaling co-operate in mediating adhesion and chemoresistance in small cell lung cancer (SCLC) cells.
Oncogene, **24**, 4462-4471 (2005)
152. Niida A., Oishi S., Sasaki Y., Mizumoto M., Tamamura H., Fujii N., Otaka A.
Facile access to (\mathcal{D})-alkene-containing diketopiperazine mimetics utilizing organocopper-mediated anti-S_N2' reactions.
Tetrahedron Lett. **46**(24) 4183-4186 (2005)
153. Burger M., Hartmann T., Krome M., Rawluk J., Tamamura H., Fujii N., Kipps T.J., Burger J.A.
Small peptide inhibitors of the CXCR4 chemokine receptor (CD184) antagonize the activation, migration, and antiapoptotic responses of CXCL12 in chronic lymphocytic leukemia B cells.
Blood, **106**(5) 1824-1830 (2005)
154. Tamamura H., Araki T., Ueda S., Wang Z., Oishi S., Esaka A., Trent J.O., Nakashima H., Yamamoto N., Peiper S.C., Otaka A., Fujii N.
Identification of novel low molecular weight CXCR4 antagonists by structural tuning of cyclic tetrapeptide scaffolds.
J. Med. Chem. **48**(9) 3280-3289 (2005)
155. Bhonsle J.B., Wang Z., Tamamura H., Fujii N., Peiper S.C., Trent J.O.
A simple, automated quasi-4D-QSAR, quasi-multi way PLS approach to develop highly predictive QSAR Models for highly flexible CXCR4 inhibitor cyclic peptide ligands using scripted common molecular modeling tools.

QSAR Comb. Sci. **24**(5) 620-630 (2005)

156. Ichikawa N., Kasai S., Suzuki N., Nishi N., Oishi S., Fujii N., Kadoya Y., Hatori K., Mizuno Y., Nomizu M., Arikawa-Hirasawa E.
Identification of neurite outgrowth active sites on the laminin α 4 chain G domain.
Biochemistry, **44**(15) 5755-5762 (2005)
157. Percherancier Y., Berchiche Y., Slight I., Volkmer-Engert R., Tamamura H., Fujii N., Bouvier M., Heveker N.
Bioluminescence resonance energy transfer reveals ligand-induced conformational changes in CXCR4 homo- and heterodimers.
J. Biol. Chem. **280**(11) 9895-9903 (2005)
158. Nameki D., Kodama E., Ikeuchi M., Mabuchi N., Otaka A., Tamamura H., Ohno M., Fujii N., Matsuoka M.
Mutations conferring resistance to human immunodeficiency virus type 1 fusion inhibitors are restricted by gp41 and rev-responsive element functions.
J. Virol. **79**(2) 764-770 (2005)
159. Zannettino A.C., Farrugia A.N., Kortesidis A., Manavis J., To L.B., Martin S.K., Diamond P., Tamamura H., Lapidot T., Fujii N., Gronthos S.
Elevated serum levels of stromal-derived factor-1 α are associated with increased osteoclast activity and osteolytic bone disease in multiple myeloma patients.
Cancer Res. **65** 1700-1709 (2005)
160. Tamamura H., Hiramatsu K., Ueda S., Wang Z., Kusano S., Terakubo S., Trent J.O., Peiper SC., Yamamoto N., Nakashima H., Otaka A., Fujii N.
Stereoselective synthesis of [L-Arg-L/D-3-(2-naphthyl)alanine]-type (*E*)-alkene dipeptide isosteres and its application to the synthesis and biological evaluation of pseudopeptide analogues of the CXCR4 antagonist FC131.
J. Med. Chem. **48**(2) 380-391 (2005)
161. Piovan E., Tosello V., Indraccolo S., Cabrelle A., Baesso I., Trentin L., Zamarchi R., Tamamura H., Fujii N., Semenzato G., Chieco-Bianchi L., Amadori A.
Chemokine receptor expression in EBV-associated lymphoproliferation in hu/SCID mice: implications for CXCL12/CXCR4 axis in lymphoma generation.
Blood, **105**(3) 931-939 (2005)
162. Allen C.D., Ansel K.M., Low C., Lesley R., Tamamura H., Fujii N., Cyster J.G.
Germinal center dark and light zone organization is mediated by CXCR4 and CXCR5.

Nat. Immunol. **5** 943-952 (2004)

163. Yokoyama F., Suzuki N., Haruki M., Nishi N., Oishi S., Fujii N., Utani A., Kleinman H.K., Nomizu M.

Cyclic peptides from the loop region of the laminin alpha4 chain LG4 module show enhanced biological activity over linear peptides.

Biochemistry, **43**(42) 13590-13597 (2004)

164. Otaka A., Ueda S., Tomita K., Yano Y., Tamamura H., Matsuzaki K., Fujii N.
Facile synthesis of membrane-embedded peptides utilizing lipid bilayer-assisted chemical ligation.

Chem. Commun. **7**(15) 1722-1723 (2004)

165. Yamamoto N., Yang R., Yoshinaka Y., Amari S., Nakano T., Cinatl J., Rabenau H., Doerr H.W., Hunsmann G., Otaka A., Tamamura H., Fujii N., Yamamoto N.
HIV protease inhibitor, nelfinavir, inhibits replication of SARS-associated coronavirus.

Biochem. Biophys. Res. Commun. **318**(3) 719-725 (2004)

166. Tamamura H., Fujisawa M., Hiramatsu K., Mizumoto M., Nakashima H., Yamamoto N., Otaka A., Fujii N.

Identification of a CXCR4 antagonist, a T140 analog, as an anti-rheumatoid arthritis agent.

FEBS Lett. **569**(1-3) 99-104 (2004)

167. Peng H., Huang Y., Rose J., Erichsen D., Herek S., Fujii N., Tamamura H., Zheng J.

Stromal cell-derived factor 1 mediated CXCR4 signaling in rat and human cortical neural progenitor cells.

Journal of Neuroscience Research, **76**(1) 35-50 (2004)

168. Takenaga M., Tamamura H., Hiramatsu K., Nakamura N., Yamaguchi N., Kitagawa A., Kawai S., Mizushima Y., Nakashima H., Fujii N., Igarashi R.

A single treatment with microcapsules containing a CXCR4 antagonist suppressed pulmonary metastasis of murine melanoma.

Biochem. Biophys. Res. Commun. **320**(1) 226-232 (2004)

169. Otaka A., Watanabe J., Yukimasa A., Sasak Y., Watanabe H., Kinoshita T., Oishi S., Tamamura H., Fujii N.

SmI₂-mediated reduction of γ, γ -difluoro- α, β -enoates and its application to the synthesis of (\mathcal{D})-fluoroalkene-type dipeptide isosteres.

J. Org. Chem. **69**(5) 1634-1645 (2004)

170. Mori T., Doi R., Masui T., Koizumi M., Toyoda E., Tulachan S.S., Ito D., Kami

- K., Masui T., Fujimoto K., Tamamura H., Hiramatsu K., Fujii N., Imamura M. CXCR4 antagonist inhibits stromal cell-derived factor 1-induced migration and invasion of human pancreatic cancer.
Molecular Cancer Therapeutics, **3** 29-37 (2004)
171. Masui T., Dori R., Mori T., Tsuji S., Nakajima S., Koizumi M., Toyoda E., Tulachan S.S., Ito D., Kami K., Kuhara T., Fujii N., Oishi S., Niida A., Peiper S.C., Broach J.R., Imamura M.
Metastin and its variant forms suppress migration of pancreatic cancer cells.
Biochem. Biophys. Res. Commun. **315**(1) 85-92 (2004)
172. Ohno H., Takemoto Y., Fujii N., Tanaka T., Ibuka T.
Stereodivergent synthesis of chiral 2-alkenylaziridines: palladium(0)-catalyzed 2,3-*cis*-selective aziridination and base-mediated 2,3-*trans*-selective aziridination.
Chem. Pharm. Bull. **52**(1) 111-119 (2004)
173. Yanada R., Koh Y., Nashimori N., Matsumura A., Obika S., Mitsuya H., Fujii N., Takemoto Y.
Indium-mediated atom-transfer and reductive radical cyclizations of iodoalkynes: synthesis and biological evaluation of HIV-protease inhibitors.
J. Org. Chem. **69**(7) 2417-2422 (2004)
174. Tamamura H., Mizumoto M., Hiramatsu K., Kusano S., Terakuba S., Yamamoto N., Trent J.O., Wang Z., Peiper S.C., Nakashima H., Otaka A., Fujii N.
Topochemical exploration of potent compounds using retro-enantiomer libraries of cyclic pentapeptides.
Org. Biomol. Chem. **2**(8) 1255-1257 (2004)
175. Tamamura H., Koh Y., Ueda S., Sasaki Y., Yamasaki T., Aoki M., Maeda K., Watai Y., Arikuni H., Otaka A., Mitsuya H., Fujii N.
Reduction of peptide character of HIV protease inhibitors that exhibit nanomolar potency against multi-drug resistant HIV-1 strains.
J. Med. Chem. **46**(9) 1764-1768 (2003)
176. Ogawa M., Hatano K., Oishi S., Kawasumi Y., Fujii N., Kawaguchi M., Doi R., Imamura M., Yamamoto M., Ajito K., Mukai T., Saji H., Ito K.
Direct electrophilic radiofluorination of a cyclic RGD peptide for in vivo $\alpha v \beta 3$ integrin related tumor imaging.
Nucl. Med. Biol. **30**(1) 1-9 (2003)
177. Yasuda T., Poole A.R., Shimizu M., Nakagawa T., Julovi S.M., Tamamura H., Fujii N., Nakamura T.

- Involvement of CD44 in induction of matrix metalloproteinases by a COOH-terminal heparin-binding fragment of fibronectin in human articular cartilage in culture.
Arthritis & Rheumatism, **48**(5) 1271-1280 (2003)
178. Fujii N., Oishi S., Hiramatsu K., Araki T., Ueda S., Tamamura H., Otaka A., Kusano S., Terakubo S., Nakashima H., Broach J.A., Trent J.H., Wang X.Z., Peiper S.C.
Molecular size reduction of a potent CXCR4-chemokine antagonists using orthogonal combination of conformation-based and sequence-based libraries.
Angew. Chem. Int. Ed. Engl. **42**(28) 3251-3253 (2003)
179. Tamamura H., Hori A., Kanzaki N., Hiramatsu K., Mizumoto M., Nakashima H., Yamamoto N., Otaka A., Fujii N.
T140 analogs as CXCR4 antagonists identified as anti-metastatic agents in the treatment of breast cancer.
FEBS Lett. **550**(1-3) 79-83 (2003)
180. Otaka A., Yukimasa A., Watanabe J., Sasaki Y., Oishi S., Tamamura H., Fujii N.
Application of samarium diiodide (SmI_2)-induced reduction of γ -acetoxy- α , β -enoates with α -specific kinetic electrophilic trapping for the synthesis of amino acid derivatives.
Chem. Commun. **15** 1834-1835 (2003)
181. Tamamura H., Kato T., Otaka A., Fujii N.
Synthesis of potent β -secretase inhibitors containing a hydroxyethylamine dipeptide isostere and their structure-activity relationship studies.
Org. Biomol. Chem. **1**(14) 2468-2473 (2003)
182. Hanna J., Wald O., Goldman-Wohl D., Pruss D., Markel G., Gazit R., Katz G., Haimov-Kochman R., Fujii N., Yagel S., Peled A., Mandelboim O.
CXCL12 expression by invasive trophoblasts induces the specific migration of CD16 negative human natural killer cells.
Blood, **102**(5) 1569-1577 (2003)
183. Trent J.O., Wang Z.X., Murray J.L., Shao W., Tamamura H., Fujii N., Peiper S.C.
Lipid bilayer simulations of CXCR4 with inverse agonists and weak partial agonists.
J. Biol. Chem. **278**(47) 47136-47144 (2003)
184. Burger M., Glodek A., Hartmann T., Schmitt-Graff A., Seilberstein L.E., Fujii N., Kipps T.J., Burger J.A.

- Functional expression of CXCR4(CD184) on small cell lung cancer cells (SCLC) mediates migration, integrin activation, and adhesion to stromal cells.
Oncogene, **22**(50) 8093-8101 (2003)
185. Tamamura H., Hiramatsu K., Kusano S., Terakubo S., Yamamoto N., Trent J.O., Wang Z.X., Peiper S.C., Nakashima H., Otaka A., Fujii N.
Synthesis of potent CXCR4 inhibitors possessing low cytotoxicity and improved biostability based on T140 derivatives.
Org. Biomol. Chem. **1**(21) 3656-3662 (2003)
186. Tamamura H., Hiramatsu K., Mizumoto M., Ueda S., Kusano S., Terakubo S., Akamatsu M., Yamamoto N., Trent J.O., Wang Z.X., Peiper S.C., Nakashima H., Otaka A., Fujii N.
Enhancement of the T140-based pharmacophores leads to the development of more potent and bio-stable CXCR4 antagonists.
Org. Biomol. Chem. **1**(21) 3663-3669 (2003)
187. Oonuma T., Morimatsu M., Nakagawa T., Uyama R., Sasaki N., Tamamura H., Fujii N., Hashimoto S., Yamamura H., Syuto B.
Role of CXCR4 and SDF-1 in mammary tumor metastasis in the cat.
J. Vet. Med. Sci. **65**(10) 1069-1073 (2003)
188. Suzuki N., Nakatsuka H., Mochizuki M., Nishi N., Kadoya Y., Utani A., Oishi S., Fujii N., Kleinman H.K., Nomizu M.
Biological activities of homologous loop regions in the laminin alpha chain G domains.
J. Biol. Chem. **278**(46) 45697-45705 (2003)
189. Ogawa M., Hatano K., Oishi S., Kawasumi Y., Fujii N., Kawaguchi M., Doi R., Imamura M., Yamamoto M., Ajito K., Mukae T., Saji H., Ito K.
Direct electrophilic radiofluorination of a cyclic RGD peptide for in vivo avb3 integrin related tumor imaging.
Nucl. Med. Biol. **30**(1) 1-9 (2003)
190. Tamamura H., Kato T., Otaka A., Fujii N.
Synthesis of potent β -secretase inhibitors-containing a hydroxyethylamine dipeptide isostere and their structure-activity relationship studies.
Organic & Biomolecular Chemistry, **1**(14) 2468-2473 (2003)
191. Ara T., Itoi M., Kawabata K., Egawa T., Tokoyoda K., Sugiyama T., Fujii N., Amagai T., Nagasawa T.
A Role of CXC Chemokine Ligand 12/Stromal Cell-Derived Factor-1/Pre-B Cell Growth Stimulating Factor and Its Receptor CXCR4 in Fetal and Adult T Cell

- Development in Vivo.
Journal of Immunology, **170**(9) 4649-4655 (2002)
192. Ohno H., Miyaura K., Tanaka T., Oishi S., Toda A., Takemoto Y., Fujii N., Ibuka T.
Synthesis of allenes from allylic alcohol derivatives bearing a bromine atom using a palladium(0)/diethylzinc system.
J. Org. Chem. **67**(4) 1359-1367 (2002)
193. Tamamura H., Hori T., Otaka A., Fujii N.
Efficient stereoselective synthesis of peptidomimetics containing hydroxyethylamine dipeptide isosteres utilizing the aza-Payne rearrangement and O,N-acyl transfer reactions.
J. Chem. Soc. Perkin Trans I, 577-580 (2002)
194. Tamamura H., Hiramatsu K., Miyamaoto K., Omagari A., Oishi S., Nakashima H., Yamamoto N., Kuroda K., Nakagawa T., Otaka A., Fujii N.
Synthesis and evaluation of pseudopeptide analogs of a specific CXCR4 inhibitor, T140: the insertion of an (E)-alkene dipeptide isostere into the β II'-turn moiety.
Bioorg. Med. Chem. Lett. **12**(6) 923-928 (2002)
195. Zhang W.-bo., Navenot J.-M., Bodduluri H., Tamamura H., Hiramatu K., Omagari A., Pei G., Manfredi J.P., Fujii N., Broach J., Peiper S.C.
A point mutation that confers constitutive activity to CXCR4 reveals T140 is an inverse agonist and AMD3100 and ALX40-4C are weak partial agonists.
J. Biol. Chem. **277**(27) 24515-24521 (2002)
196. Oishi S., Kamano T., Niida A., Odagaki Y., Tamamura H., Otaka A., Hamanaka N., Fujii N.
Diastereoselective synthesis of ϕ [(E)-CH=CMe]- and ϕ [(Z)-CH=CMe]-type dipeptide isosteres by organocopper-mediated anti-SN_{2'} reaction.
Org. Lett. **4**(7) 1051-1054 (2002)
197. Oishi S., Niida A., Kamano T., Odagaki Y., Tamamura H., Otaka A., Hamanaka N., Fujii N.
Diastereoselective synthesis of ϕ [(E)-CMe=CH]- and ϕ [(E)-CMe=CMe]-type dipeptide isosteres based on organocopper-mediated anti-SN_{2'} reaction.
Org. Lett. **4**(7) 1055-1058 (2002)
198. Anzai M., Yanada R., Fujii N., Ohno H., Ibuka T., Takemoto Y.
Asymmetric synthesis of β 2,3-amino acids by InI-Pd(0)-promoted metalation and addition of chiral 2-vinylaziridines.
Tetrahedron, **58**(26) 5231-5239 (2002)

199. Yanada R., Nishimori N., Matsumura A., Fujii N., Takemoto Y.
Indium-mediated atom-transfer cyclization and reductive cyclization.
Tetrahedron Lett. **43**(26) 4585-4588 (2002)
200. Otaka A., Nakamura M., Nameki D., Kodama E., Uchiyama S., Nakamura S.,
Nakano H., Tamamura H., Kobayashi Y., Matsuoka M., Fujii N.
Remodeling of gp41-C34 peptide leads to highly effective inhibitors of the fusion
of HIV-1 with target cells.
Angew. Chem. Int. Ed. Engl. **41**(16) 2937-2940 (2002)
201. Tamamura H., Omagari A., Hiramatsu K., Oishi S., Habashita H., Kanamoto T.,
Gotoh K., Yamamoto N., Nakashima H., Otaka A., Fujii N.
Certification of the critical importance of L-3-(2-naphthyl)alanine at position 3 of
a specific CXCR4 inhibitor, T140, leads to an exploratory performance of its
downsizing study.
Bioorg. Med. Chem. **10**(5) 1417-1426 (2002)
202. Hosotani R., Miyamoto Y., Fujimoto K., Doi R., Otaka A., Fujii N., Imamura M.
Trojan p16 peptide suppresses pancreatic cancer growth and prolongs survival
in mice.
Clin. Cancer Res. **8**(4) 1271-1276 (2002)
203. Oishi S., Niida A., Kamano T., Miwa Y., Taga T., Odagaki Y., Hamanaka N.,
Yamamoto M., Akaji K., Tamamura H., Otaka A., Fujii N.
Regio- and stereoselective ring-opening of chiral 1,3-oxazolidin-2-one derivatives
by organocopper reagents provides novel access to di-, tri- and tetra-substituted
alkene dipeptide isosteres.
J. Chem. Soc. Perkin Trans I, 1786-1793(2002)
204. Otaka A., Katagiri F., Kinoshita T., Odagaki Y., Oishi S., Tamamura H.,
Hamanaka N., Fujii N.
Regio- and stereoselective synthesis of (E)-alkene trans-Xaa-Pro dipeptide
mimetics utilizing organocopper-mediated anti-SN2' reactions.
J. Org. Chem. **67**(17) 6152-6161 (2002)
205. Oishi S., Kamano T., Niida A., Odagaki Y., Hamanaka N., Yamamoto M., Ajito
K., Tamamura H., Otaka A., Fujii N.
Diastereoselective synthesis of new psi[(E)-CH=CMe]- and
psi[(Z)-CH=CMe]-type alkene dipeptide isosteres by organocopper reagents and
application to conformationally restricted cyclic RGD peptidomimetics.
J. Org. Chem. **67**(17) 6162-6173 (2002)
206. Lee J.U., Hosotani R., Wada M., Doi R., Koshiba T., Fujimoto K., Miyamoto Y.,

- Tsuji S., Nakajima S., Hirohashi M., Uehara T., Arano Y., Fujii N., Imamura M. Antiproliferative activity induced by the somatostatin analogue, TT-232, in human pancreatic cancer cells.
Eur. J. Cancer **38**(11) 1526-1534 (2002)
207. Owen S.M., Rudolph D., Schols D., Fujii N., Yamamoto N., Lal R.B. Susceptibility of diverse primary HIV isolates with varying co-receptor specificity's to CXCR4 antagonistic compounds.
J. Med. Virol. **68**(2) 147-155 (2002)
208. Petit I., Szypers-Kravitz M., Nagler A., Lahav M., Peled A., Habler L., Ponomaryov T., Taichman R., Arenzana-Seisdedos F., Fujii N., Sandbank J., Zipori D., Lapidot T. G-CSF induces stem cell mobilization by decreasing bone marrow SDF-1 and up-regulating CXCR4.
Nature Immunology, **3** 687-694 (2002)
209. Zhang W.-bo., Navenot J.-M., Bodduluri H., Tamamura H., Hiramatu K., Omagari A., Pei G., Manfredi J.P., Fujii N., Broach J., Peiper S.C. CXCR4 variants with autonomous signaling conferred by a single mutation in the third transmembrane domain reveal T140 is an inverse agonist and AMD3100 and ALX40-4C are weak partial agonists.
J. Biol. Chem. **277**(27) 24515-24521 (2002)
210. K. Gotoh, M. Yoshimori, K. Kanbara, H. Tamamura, T. Kanamoto, K. Mochizuki, N. Fujii & H. Nakashima Increase of R5 HIV-1 Infection and CCR5 Expression in T cells Treated with High Concentration of CXCR4 Antagonists and SDF-1.
J. Infecti. chemotherapy., **7**, 28-36 (2001)
211. A. Otaka, H. Watanabe, E. Mtsuyama, A. Yukimasa, H. Tamamura & N. Fujii Synthesis of (\mathcal{D})-Fluoroalkene Dipeptide Isosteres Utilizing Organocopper-mediated Reduction of γ,γ -difluoro- α,β -enoates.
Tetrahedron Lett., **42**, 285-287 (2001)
212. H. Tamaura, M. Sugioka, Y. Odagaki, A. Omagari, Y. Kan, S. Oishi, H. Nakashima, N. Yamamoto, S. C. Peiper, N. Hamanaka, A. Otaka & N. Fujii Conformational Study of a Highly Specific CXCR4 Inhibitor, T140, Disclosing the Close Proximity of Its Intrinsic Pharmacophores Associated with Strong Anti-HIV Activity .
Bioorg. Med. Chem. Lett., **11**, 359-362 (2001)
213. K. Kanbara, S. Sato, J. Tanuma, H. Tamamura, K. Gotoh, M. Yoshimori, T.

- Kanamoto, M. Kitano, N. Fujii & H. Nakashima
Biological and Genetic Characterization of a Human Immunodeficiency Virus Strain Resistant to CXCR4 Antagonist T134.
Aids Res. Human Retroviruses., **17**, 615-622 (2001)
214. Y. Takemoto, M. Anzai, R. Yanada, N. Fujii, H. Ohno & T. Ibuka
Stereoselective Synthesis of Nonracemic 1,3-Amino Alcohols from Chiral 2-vinylaziridines by In(I)-Pd(0)-promoted Metalation.
Tetrahedron Lett., **42**, 1725-1728 (2001)
215. H. Tamamura, A. Omagari, K. Hiramatsu, K. Gotoh, T. Kanamoto, Y. Xu, E. Kodama, M. Matsuoka, T. Hattori, N. Fujii, H. Nakashima, A. Otaka & N. Fujii
Development of Specific CXCR4 Inhibitors Possessing High Selectivity Indexes as Well as Complete Stability in Serum Based on an Anti-HIV Peptide T140.
Bioorg. Med. Chem. Lett., **11**, 1897-1902 (2001)
216. T. Egawa, K. Kawabata, H. Kawamoto, K. Amada, R. Okamoto, N. Fujii, T. Kishimoto, Y. Katsura & T. Nagasawa
The Earliest Stages of B Cell Development Require a Chemokine Stronal Cell-Derived Factor SDF-1/Pre-B Cell Growth-Stimulating Factor.
Immunity, **15**, 323-334 (2001)
217. H. Tamamura, A. Omagari, K. Hiramatsu, T. Kanamoto, K. Gotoh, K. Kanbara, N. Yamamoto, H. Nakashima, A. Otaka & N. Fujii
Synthesis and Evaluation of Bifunctional Anti-HIV Agents and Specific CXCR4 Antagonists-AZT Conjugation.
Bioorg. Med. Chem., **9**, 2179-2187 (2001)
218. H. Ohno, M. Anzai, A. Toda, S. Ohishi, N. Fujii, T. Tanaka, Y. Takemoto & T. Ibuka
Stereoselective Synthesis of 2-Alkenylaziridines and 2-Alkenylazetidines by Palladium-Catalyzed Intramolecular Amination of α - and β -Amino Allenes
J. Org. Chem., **66**, 4904 -4914 (2001)
219. A. Otaka, H. Watabe, A. Yukimasa, S. Oishi, H. Tamamura & N. Fujii
New Access to α -Substituted (\mathcal{D} -Fluoroalkene Dipeptide Isosteres Utilizing Organocopper Reagents under 'Reduction-Oxidative Alkylation (R-OA)' Conditions.
Tetrahedron Lett., **42**, 5443-5446 (2001)
220. S. Oishi, H. Tamamura, M. Yamashita, A. Otaka & N. Fujii
Stereoselective synthesis of a set of two functionalized (E)-alkene dipeptide isosteres of L-amino acid-L Glu and L-amino acid-D-Glu.

J. Chem. Soc., Perkin Trans I, 2001, 2445-2451

221. T. Hara, Y. Mitani, K. Tanaka, N. Uematsu, A. Takakura, T. Tachi, H. Kodama, M. Kondo, H. Mori, A. Otaka, N. Fujii, & K. Matsuzaki
Heterodimer Formation between the Antimicrobial Peptides Magainin 2 and PGLa in Lipid Bilayers: A Cross-Linking Study.
Biochemistry, **40**(41), 12395-12399 (2001)
222. T. Tanaka, A. Yoshida, E. Baba, T. Murakami, J. Lichtenfeld, T. Omori, T. Kimura, N. Tsurutani, N. Fujii, Zi-X. Wang, S.C. Peiper, N. Yamamoto & Y. Tanaka
A Unique Monoclonal Antibody Recognizing the Third Extracellular loop of the Human CXCR4 Induces Lymphocyte Agglutination and Enhances Human Immunodeficiency Virus Type-1-Mediated Syncytium Formation and Productive infection.
J. Virol. **75** (23), 11534-11543 (2001)
223. M. Kawaguchi, R. Hosotani, S. Oishi, N. Fujii, S. Tulachan, M. Koizumi, E. Toyoda, T. Masui, S. Nakajima, S. Tsuji, J. Ida, K. Fujimoto, M. Wada, R. Doi and M. Imamura
A novel synthetic Arg-Gly-Asp containing Peptide cyclo(-RGDf=V) is the potent inhibitor of angiogenesis.
Biochem. Biophys. Res. Commun., **288**, 711-717 (2001)
224. K. Hodohara, N. Fujii, N. Yamamoto & K. Kaushansky
Stromal cell-derived factor-1 (SDF-1) acts together with thrombopoietin to enhance the development of megakaryocytic progenitor cells (CFU-MK).
Blood, **95**(3), 769-775 (2000)
225. H. Ohno, A. Toda, N. Fujii, Y. Takemoto, T. Tanaka & T. Ibuka
Stereoselective Synthesis of Chiral Amino Allenes by Organocopper-Mediated anti-SN2'-Substitution Reaction of Chiral Ethynylaziridines.
Tetrahedron, **56**, 2811-2820 (2000)
226. Otaka, E. Mitsuyama, T. Kinoshita, H. Tamamura & N. Fujii
Stereoselective Synthesis of CF2-Substituted Phosphothreonine Mimetics and Their Incorporation into Peptides Using Newly Developed Deprotection Procedures.
J. Org. Chem., **65**(16), 4888-4899 (2000)
227. K. Miyoshi, A. Otaka, M. Kaneko, H. Tamamura & N. Fujii
A New Practical Strategy for the Synthesis of Long Chain Phosphopeptide.
Chem. Pharm. Bull., **48**(8) 1230-1233 (2000)

228. A. Otaka, E. Mitsuyama, H. Watanabe, H. Tamamura & N. Fujii
Development of New Methodology for the Synthesis of Functionalized
 α -Fluoro-phosphonates and Its Practical Application to the Preparation of
Phosphopeptide Mimetics.
Chem. Commun., **2000**, 1081-1082
229. H. Ohno, A. Toda, S. Oishi, T. Tanaka, Y. Takemoto, N. Fujii & T. Ibuka
Novel Synthesis of Chiral Terminal Allenes via Palladium(0)-catalyzed
Reduction of Mesylates of 2-Bromobut-2-en-1-ols Bearing a Protected Amino
Group, Using Diethylzinc
Tetrahedron Lett., **41**, 5131-5134 (2000)
230. Koshiba, R. Hosotani, Y. Miyamoto, J. Ida, S. Tsuji, S. Nakamura, M.
Kawaguchi, H. Kobayashi, R. Doi, T. Hori, N. Fujii & M. Imamura
Expression of Stromal Cell-derived Factor 1 and CXCR4 Ligand Receptor
System in Pancreatic Cancer: A Possible Role for Tumor Progression.
Clin. Cancer Res., **6**, 3530-3535 (2000)
231. K. Fujimoto, R. Hosotani, Y. Miyamoto, R. Doi, T. Koshiba, A. Otaka, N. Fujii,
R.D. Beauchamp & M. Imamura
Inhibition of pRb Phosphorylation and Cell Cycle Progression by an
Antennapedia-P16(INK4A) Fusion Peptide in Pancreatic Cancer Cells.
Cancer Lett., **159**(2), 151-158 (2000)
232. K. Kanbara, N. Fujii & H. Nakashima
A Study of Anti-HIV Compounds Which Interfere the Virus Entry via Coreceptor
CXCR4
Kansenshogaku Zasshi., **74**, 237-244 (2000)
233. Y. Kuroda, K. Miyamoto, K. Tanaka, Y. Maeda, J. Ishikawa, R. Hinata, A.
Otaka, N. Fujii & T. Nakagawa
Interaction between Local Anesthetics and Na⁺-Channel Inactivation Gate
Peptides in Phosphatidylserine Suspensions as Studied by ¹H-NMR
Spectroscopy.
Chem. Pharm. Bull., **48**, 1293-1298 (2000)
234. Y. Kuroda, K. Miyamoto, M. Matsumoto, Y. Maeda, K. Kanaori, A. Otaka, N.
Fujii & T. Nakagawa
Structural Study of the Sodium Channel Inactivation Gate Peptide Including an
Isoleucine-Phenylalanine-Methionine Motif and Its Analogous Peptide
(Phenylalanine/Glutamine) in Trifluoroethanol Solutions and SDS Micelles.
J. Pept. Res., **56**, 172-184 (2000)

235. Ponomaryov, A. Peled, I. Petit, R. Taichman, L. Habler, J. Sandbank, F.A. Seidsdedos, A. Magerus, A. Caruz, N. Fujii, A. Nagler, M. Lahav, M. Szyper-Kravit, D. Zipori & T. Lapidot
Induction of the Chemokine Stromal-derived Factor-1 Following DNA Damage Improves Human Stem Cell Function.
J. Clin. Invest., **106**, 1331-1339 (2000)
236. H. Tamamura, A. Omagari, S. Oishi, T. Kanamoto, N. Yamamoto, S. C. Peiper, H. Nakashima, A. Otaka & N. Fujii
Pharmacophore Identification of a Specific CXCR4 Inhibitor, T140, Leads to Development of Effective Anti-HIV Agents with Very High Selectivity Indexes
Bioorg. Med. Chem. Lett., **10**, 2633-2637 (2000)
237. H. Ohno, A. Toda, Y. Miwa, T. Taga, N. Fujii & T. Ibuka
Synthesis of Chiral Amino Allenes via an Organocyanocupurate-Mediated Ring-Opening Reaction of Enantiopure Ethynylaziridines.
Tetrahedron Lett., **40**, 349-352 (1999)
238. Arakaki, H. Tamamura, M. Premenathan, K. Kanbara, S. Ramanan, K. Mochizuki, M. Baba, N. Fujii & H. Nakashima
T134, a smallmolecule CXCR4 inhibitor, has no cross drug-resistance with the different class of CXCR4 antagonist AMD3100.
J. Virol., **73**, 1719-1723 (1999).
239. Y. Xu, H. Tamamura, R. Arakaki, H. Nakashima, X. Zhang, N. Fujii, T. Uchiyama & T. Hattori
Marked Increase in Anti-HIV Activity, as Well as Inhibitory Activity against HIV Entry Mediated by CXCR4, Linked to Enhancement of the Binding Ability of Tachyplesin Analogs to CXCR4.
AIDS Research and Human Retroviruses., **15**, 419-427 (1999)
240. T. Murakami, Z-H. Lu, Y. Koyanagi, Y. Tanaka, J. Kim, Y. Suzuki, S. Minoguchi, H. Tamamura, M. Waki, A. Matsumoto, N. Fujii, H. Shida, J.A. Hoxie, S.C. Peiper & N. Yamamoto
Inhibitory Mechanism of the CXCR4 Antagonist T22 against Human Immunodeficiency Virus Type1 Infection.
J. Virol., **73**(9), 7489-7496 (1999)
241. H. Ohno, A. Toda, N. Fujii, Y. Miwa, T. Taga, Y. Yamaoka, E. Osawa & T. Ibuka
Sterically Congested Chiral Activated Aziridines: Synthesis of Both 2,3-Cis- and 2,3-Trans-2-Alkenyl-3-alkylaziridines from Common Intermediates.
Tetrahedron Lett., **40**, 1331-1334 (1999)

242. H. Ohno, A. Toda, Y. Miwa, T. Taga, E. Osawa, Y. Yamanaka, N. Fujii & T. Ibuka
First Palladium-Catalyzed Aziridination Reaction of Amino Allenes.
J. Org. Chem., **64** (9), 2992-2993 (1999)
243. K. Ishii, H. Ohno, Y. Takemoto, E. Osawa, Y. Yamaoka, N. Fujii & T. Ibuka
Selective Synthesis of *cis*-2-Vinyl-3-Alkylaziridines and 3-Pyrrolines from Common Intermediates (Z -4-*N*-Arylsulfonylaminoalk-2-en-1-ols.
J. Chem. Soc., Perkin Trans. I., 2155-2163 (1999)
244. Y. Kuroda, Y. Maeda, K. Miyamoto, K. Tanaka, K. Kanaori, A. Otaka, N. Fujii, & T. Nakagawa
1H-NMR and Circular Dichroism Spectroscopic Studies on Changes in Secondary Structures of the Sodium Channel Inactivation Gate Peptides as Caused by the Pentapeptide KIFMK
Biophysical J., **77**, 1363-1373 (1999)
245. H. Ohno, A. Toda, Y. Takemoto, N. Fujii and T. Ibuka
Convenient Syntheses of Chiral 3-Substituted 2-Ethynylaziridines.
J. Chem. Soc., Perkin Trans. I., **1999**, 2949-2962.
246. H. Tamamura, M. Yamashita, Y. Nakajima, K. Sakano, A. Otaka, H. Ohno, T. Ibuka & N. Fujii
Regiospecific Ring-Opening Reactions of β -Aziridinyl α,β -Enoates with Acids: Application to the Stereoselective Synthesis of a Couple of Diastereoisomeric (*E*)-Alkene Dipeptide Isosteres from a Single β -Aziridinyl α,β -Enoate and to the Convenient Preparation of Amino Alcohols Bearing α,β -Unsaturated Ester Groups.
J. Chem. Soc., Perkin Trans I., **1999**, 2983-2996.
247. M. Anzai, A. Toda, H. Ohno, Y. Takemoto, N. Fujii & T. Ibuka
Palladium-catalyzed regio- and stereoselective synthesis of *N*-protected 2,4-dialkylated azacyclobutanes from amino allenes.
Tetrahedron Lett., **40**, 7393-7397 (1999)
248. P. Buffet, B. Gamain, C. Sheiding, D. Baruch, J. Smith, R. Hernandez-Rivas, B. Pouvelle, S. Oishi, N. Fujii, T. Fusai, D. Parzy, L. Miller & A. Scherf
Plasmodium Falciparum Domain Mediating Adhesion to Chondroitin Sulfate A: A Receptor for Human Placental Infection.
Proc. Natl. Acad. Sci., USA., **96** (22), 12743-12748 (1999)
249. Uehara, Y. Arano, M. Ono, Y. Fujioka, K. Ogawa, S. Namba, M. Nakayama, M. Koizumi, N. Fujii, K. Horiuchi, A. Yokoyama & H. Saji

The Integrity of the Disulfide Bond in a Cyclic Somatostatin Analog during ^{99m}Tc Complexation Reactions.

Nuclear Medicine & Biology, **26**, 883-890 (1999)

250. H. Tamamura, R. Arakaki, H. Funakoshi, M. Imai, A. Otaka, T. Ibuka, H. Nakashima, T. Murakami, M. Waki, A. Matsumoto, N. Yamamoto & N. Fujii Effective Lowly Cytotoxic Analogs of an HIV-Cell Fusion Inhibitor, T22 ([Tyr_{5,12},Lys₇]-Polyphemusin II).
Bioorg. Med. Chem., **6**, 231-238 (1998)
251. H. Tamamura, T. Ishihara, H. Oyake, M. Imai, A. Otaka, T. Ibuka, R. Arakaki, H. Nakashima, T. Murakami, M. Waki, A. Matsumoto, N. Yamamoto & N. Fujii Convenient One-Pot Synthesis of Cystine-Containing Peptides Using the Trimethylsilyl Chloride-Dimethylsulfoxide/Trifluoroacetic Acid System and Its Application to the Synthesis of Bifunctional Anti-HIV Compoundds.
J. Chem. Soc., Perkin Trans. I, **1998**, 495-500 (1998)
252. H. Tamamura, F. Matsumoto, K. Sakano, A. Otaka, T. Ibuka & N. Fujii Unambiguous Synthesis of Stromal Cell-Derived Factor-1 by Regioselective Disulfide Bond Formation Using a Dimethylsulfoxide/ Aqueous HCl System.
J. Chem. Soc., Chem. Commun., **1998**, 151-152 (1998)
253. H. Uenishi, N. Iwatani, K. Kuribayashi, H. Tamamura, N. Fujii, T. Nakatani, T. Kawasaki & H. Yamagishi Overlapping Epitopes of Friend Murine Leukemia Virus *gag*-Encoded Leader Sequence Recognized by Single Cytotoxic T-Lymphocyte Clones.
Immunol. Lett., **62**, 33-38 (1998)
254. H. Uenishi, N. Iwanami, H. Yamagishi, T. Nakatani, T. Kawasaki, H. Tamamura, N. Fujii & K. Kuribayashi Induction of Crossreactivity in an Endogenous Viral Peptide Non-Reactive to FBL-3 Tumor-Specific Helper T Cell Clones.
Microbial Immunol., **42**(7), 479-484 (1998)
255. H. Habashita, T. Kawasaki, Y. Takemoto, N. Fujii & T. Ibuka A New Synthesis of the Central Substructure of Botryococcenes.
J. Org. Chem., **63** (7), 2392-2396 (1998).
256. H. Tamamura, M. Waki, M. Imai, A. Otaka, T. Ibuka, K. Waki, K. Miyamoto, A. Matsumoto, T. Murakami, H. Nakashima, N. Yamamoto & N. Fujii Downsizing of an Anti-HIV Peptide, T22 ([Tyr_{5,12},Lys₇]-Polyphemusin II), with Maintenance of Anti-HIV Activity and Solution Structure.

Bioorg. Med. Chem., **6**, 473-479 (1998)

257. H. Tamamura, T. Ishihara, M. Masuda, H. Funakoshi, H. Oyake, M. Imai, T. Murakami, R. Arakaki, H. Nakashima, A. Otaka, T. Ibuka, M. Waki, A. Matsumoto, N. Yamamoto & N. Fujii
Pharmacophore Identification of a Chemokine Receptor (CXCR4) Antagonist, T22 ([Tyr5,12, Lys7]-Polyphemusin II) which Specifically Blocks T Cell-line-tropic HIV-1 Infection.
Bioorg. Med. Chem., **6**, 1033-1041 (1998)
258. Toda, H. Aoyama, N. Mimura, H. Ohno, N. Fujii & T. Ibuka
Reactions of *N*-Arylsulfonyl-2,3-*cis*-and *N*-Arylsulfonyl-2,3-*trans*-3-alkyl-2-2-vinylaziridines with Organocopper Reagents: Importance of 2,3-*cis*-Stereochemistry in Controlling Regio- and Stereoselectivity.
J. Org. Chem., **63** (20), 7053-7061 (1998)
259. T. Hattori, H. Komoda, S. Pahwa, M. Tateyama, X. Zhang, Y. Xu, S. Ogawa, H. Tamamura, N. Fujii, K. Fukutake & T. Uchiyama
Decline of anti-DP107 Antibody Associated with Clinical Progression.
AIDS, **12** (12), 1557-1559 (1998)
260. H. Ohno, K. Ishii, A. Honda, H. Tamamura, N. Fujii, Y. Takemoto & T. Ibuka
A 2,3-*cis*-Selective Synthesis of Aziridines Bearing a Vinyl Group from Allyl Methyl Carbontes and Mesylates
J. Chem. Soc., Perkin Trans. I, **1998**, 3703-3716.
261. H. Ohno, A. Toda, N. Fujii & T. Ibuka
A Convenient Synthesis of Activated Enantiomerically Pure 2-Ethynylaziridines
Tetrahedron Assymetry, **9**, 3929-3933 (1998)
262. H. Tamamura, Y. Xu, T. Hattori, X. Zhang, R. Arakaki, K. Kanbara, A. Omagari, A. Otaka, T. Ibuka, N. Yamamoto, H. Nakashima & N. Fujii
A Low Molecular Weight Inhibitor against the Chmokine Receptor CXCR4: a Strong Anti-HIV Peptide T140.
Biochem. Biophys. Res. Commun., **253**, 877-882 (1998)
263. T. Ibuka, N. Mimura, H. Aoyama, M. Akaji, H. Ohno, Y. Miwa, T. Taga, K. Nakai, H. Tamamura & N. Fujii
A Thermodynamic Preference of Chiral *N*-Methanesulfonyl and *N*-Arenesulfonyl 2,3-*cis*-3-Alkyl-2-Vinylaziridines over Their 2,3 *trans*-Isomers: Useful Palladium (0)-Catalyzed Equilibration Reactions for the Synthesis of (*E*)-Alkene Dipeptide Isosteres.
J. Org. Chem., **62**(4), 999-1015 (1997)

264. K. Miyasaka, S. Kanai, M. Masuda, T. Ibuka, K. Nakai, N. Fujii & A. Funakoshi
Involvement of Cholinergic Processes in Cholecystokinin (CCK) Releasing by Luminal Oleic Acid.
J. Autonomic Nervous System., **63**, 179-182 (1997)
265. K. Matsuzaki, A. Nakamura, O. Murase, K. Sugishita, N. Fujii & K. Miyajima
Modulation of Magainin 2-Lipid Bilayer Interactions by Peptide Charges.
Biochemistry, **36**(6), 2104-2111 (1997)
266. K. Fujimoto, R. Doi, R. Hosotani, M. Wada, J-Uk., Lee, T. Koshiba, T. Ibuka, H. Habashita, K. Nakai, N. Fujii & M. Imamura
Effects of Structure Modification on Biological Activity of Bombesin Analogues with (*E*)-Alkene Bond.
Life Sciences., **60**(1), 29-34 (1997)
267. K. Matsuzaki, K. Sugishita, M. Harada, N. Fujii & K. Miyajima
Interaction of an Antimicrobial Peptide, Magainin 2, with Outer and Inner Membrane of Gram-Negative Bacteria.
Biochim. Biophys. Acta., **1327**, 119-130 (1997)
268. T. Ibuka, N. Mimura, H. Ohno, K. Nakai, M. Akaji, H. Habashita, H. Tamamura, Y. Miwa, T. Taga, N. Fujii & Y. Yamamoto
Palladium(0)-Catalyzed Isomerization Reactions of Aziridines Bearing an α,β -Unsaturated Ester Group: A Thermodynamic Preference for Chiral Alkyl (*2E*)-4,5-*cis*-4,5-Epimino-N-(Alkyl- or Arylsulfonyl) 2-Enoates over the Other Three Stereoisomers.
J. Org. Chem., **62** (9), 2982-2991 (1997)
269. M. Noda, T. Ibuka, H. Habashita & N. Fujii
A Highly Stereoselective Synthesis of the Functionalized (*E*)-Alkene Dipeptide Isosteres of Trp-Val via Organocyanocopper-Lewis Acid Mediated Reaction.
Chem. Pharm. Bull., **45** (8), 1259-1264 (1997)
270. T. Murakami, T. Nakajima, Y. Koyanagi, K. Tachibana, N. Fujii, H. Tamamura, N. Yoshida, M. Waki, A. Matsumoto, O. Yoshie, T. Kishimoto, N. Yamamoto & T. Nagasawa
A Small Molecule CXCR4 Inhibitor that Blocks T Cell-Line Tropic HIV-1 Infection.
J. Exp. Med., **186** (8) 1389-1393 (1997)
271. K. Matsuzaki, S. Yoneyama, N. Fujii, K. Miyajima, K. Yamada, Y. Kirino & K. Anzai

Membrane Permeabilization Mechanism of a Cyclic Antimicrobial Peptide, Tachyplesin I, and Its Linear Analogs.

Biochemistry, **36** (32), 9799-9806 (1997)

272. H. Ohno, N. Mimura, A. Otaka, H. Tamamura, N. Fujii, T. Ibuka, I. Shimizu, A. Satake & Y. Yamamoto
Palladium-Catalyzed Reductive Ring Opening with Formic Acid of Aziridines Bearing an α,β -Unsaturated Ester Group.
Tetrahedron, **53** (38), 12933-12946 (1997)
273. H. Aoyama, N. Mimura, H. Ohno, K. Ishii, A. Toda, H. Tamamura, A. Otaka, N. Fujii & T. Ibuka
Regio- and Stereoselectivity in Reactions of 2,3-*cis* and *trans*-3-Alkyl-2-vinylaziridines with Organocopper Reagents: Importance of 2,3-*cis*-Stereochemistry in Controlling Selectivity.
Tetrahedron Lett., **38**, 7383-7386 (1997)
274. H. Tamamura, M. Yamashita, H. Muramatsu, H. Ohno, T. Ibuka, A. Otaka & N. Fujii
Regiospecific Ring-Opening Reactions of Aziridines Bearing an α,β -Unsaturated Ester Group with Trifluoroacetic Acid or Methanesulfonic Acid: Application to Stereoselective Synthesis of (*E*)-Alkene Dipeptide Isosteres
Chem. Commun., **1997**, 2327-2328 (1997)
275. H. Habashita, T. Kawasaki, M. Akaji, H. Tamamura, T. Kimachi, N. Fujii & T. Ibuka
One-pot Transformation of *p*-Toluenesulfonates of 2,3-Epoxy Alcohols into Allylic Alcohols.
Tetrahedron Lett., **38** (48), 8307-8310 (1997)
276. Y. Kuroda, M. Ogawa, H. Nasu, M. Terashima, M. Kasahara, Y. Kiyama, M. Wakita, Y. Fujiwara, N. Fujii & T. Nakagawa
Locations of Local Anesthetic Dibucaine in Model Membranes and the Interaction between Dibucaine and a Na⁺-Channel Inactivation Gate Peptide as Studied by ²H- and ¹H-NMR Spectroscopies.
Biophys. J., **71** (3), 1191-1207 (1996)
277. H. Tamamura, A. Otaka, T. Murakami, T. Ishihara, T. Ibuka, M. Waki, A. Matsumoto, N. Yamamoto & N. Fujii
Interaction of an Anti-HIV peptide, T22, with GP120 and CD4.
Biochem. Biophys. Res. Commun., **219**(2), 555-559 (1996)
278. N. Fujii, H. Habashita, M. Akaji, K. Nakai, T. Ibuka, M. Fujiwara, H.

Tamamura & Y. Yamamoto

Simple One-pot Transformation of Toluene-*p*-sulfonate of 2,3-Epoxy Alcohols into Allylic Alcohols.

J. Chem. Soc., Perkin I, Comm., **1996**, 865-866 (1996)

279. N. Mimura, T. Ibuka, M. Akaji, Y. Miwa, T. Taga, K. Nakai, H. Tamamura, N. Fujii & Y. Yamamoto

An Unusual Thermodynamic Preference of Chiral *N*-Arylsulfonyl *cis*-3-Alkyl-2-vinyl-aziridines over Their *trans*-Isomers: Palladium (0)-catalyzed Equilibration Reations.

J. Chem. Soc, Chem. Comm., **1996**, 351-352 (1996)

280. T. Ibuka, M. Akaji, N. Mimura, H. Habashita, K. Nakai, H. Tamamura, N. Fujii & Y. Yamamoto

A Thermodynamic Preference of Chiral *cis*- γ,δ -Epimino-(*E*)- α,β -unsaturated Esters over Other Stereoisomers: Synthetically Useful Pd(0)-catalyzed Equilibrated Reactions of Aziridines Bearing an α,β -Unsaturated Ester Group.

Tetrahedron Lett., **37**(16), 2849-2852 (1996)

281. M. Waki, K. Waki, K. Miyamoto, A. Matsumoto, H. Tamamura, N. Fujii, T.

Murakami, H. Nakashima & N. Yamamoto

Molecular Size of an Anti-HIV Peptide, T22, Can Be Reduced without Loss of the Activity.

Chemistry Letters., 571-572 (1996)

282. H. Tamamura, T. Ishihara, T. Ibuka, A. Otaka & N. Fujii

A Convenient One-Pot Synthesis of Cystine-Containing Peptides from Protected Peptidyl Resins Using the Trimethylsilyl Chloride-Dimethylsulfoxide-Trifluoroacetic Acid System.

J. Chem. Soc., Perkin Trans I (Perkin Commun.), **1996**, 1911-1912 (1996)

283. H. Tamamura, T. Ishihara, A. Otaka, T. Murakami, T. Ibuka, M. Waki, A. Matsumoto, N. Yamamoto & N. Fujii

Analysis of the Interaction of an Anti-HIV Peptide, T22([Tyr_{5,12}, Lys₇]-Polypeptidus II), with gp 120 and CD4 by Surface Plasmon Resonance.

Biophys. Biophys. Acta., **1298**, 37-44 (1996)

284. H. Tamamura, A. Otaka, T. Murakami, T. Ibuka, K. Sakano, M. Waki, A. Matsumoto, N. Yamamoto & N. Fujii

An Anti-HIV Peptide, T22, Forms a Highly Active Complex with Zn(II).

Biochem. Biophys. Res. Commun., **229**, 648-652 (1996)

285. K. Matsuzaki, O. Murase, N. Fujii & K. Miyajima

An Antimicrobial Peptide, Magainin 2, Induced Rapid Flip-Flop of Pospholipids Coupled with Pore Formation and Peptide Translocation.

Biochemistry, **35** (35), 11361-11368 (1996)

286. T. Ibuka, K. Nakai, M. Akaji, H. Tamamura, N. Fujii & Y. Yamamoto

An Aza-Payne Rearrangement-Epoxide Ring Opening Reaction of 2-Aziridinemethanols in a One-pot Manner: A Regio- and Stereoselective Synthetic Route to Diastereomerically Pure *N*-Protected 1,2-Amino Alcohols.

Tetrahedron, **52** (36), 11739-11752 (1996)

287. H. Tamamura, A. Otaka, W. Takada, Y. Terakawa, H. Yoshizawa, M. Masuda, T. Ibuka, T. Murakami, H. Nakashima, M. Waki, A. Matsumoto, N. Yamamoto & N. Fujii

Solution Phase Synthesis of an Anti-human Immunodeficiency Virus Peptide, T22 ([Tyr_{5,12},Lys₇]-Polyphemusin II), and the Modification of Trp by the *p*-Methoxybenzyl Group of Cys during Trimethylsilyl Trifluoromethanesulfonate Deprotection.

Chem. Pharm. Bull., **43** (1), 12-18 (1995)

288. Otaka, K. Miyoshi, P. P. Roller, T. R. Burke, Jr., H. Tamamura & N. Fujii

Practical Synthesis of Phosphopeptides using Dimethyl-protected Phosphoamino Acid Derivatives.

J. Chem. Soc., Chem. Commun., **1995** (3), 387-389 (1995)

289. A. Otaka, K. Miyoshi, T.R. Burke, Jr., P.P. Roller, H. Kubota, H. Tamamura & N. Fujii

Synthesis and Application of *N*Boc-L-2-Amino-4-(diethylphosphono)-4,4-difluorobutanoic Acid for Solid Phase Synthesis of Nonhydrolyzable Phosphoserine Peptide Analogs.

Tetrahedron Lett., **36** (6), 927-930 (1995)

290. M. Wada, R. Doi, R. Hosotani, S. Higashide, T. Ibuka, H. Habashita, K. Nakai, N. Fujii & M. Imamura

Effect of a New Bombesin Receptor Antagonist, (*E*)-Alkene Bombesin Isostere, on Amylase Release from Rat Pancreatic Acini.

Pancreas, **10** (3), 301-305 (1995)

291. T. Ibuka, K. Nakai, H. Habashita, Y. Hotta, A. Otaka, H. Tamamura, N. Fujii, N. Mimura, Y. Miwa, T. Taga, Y. Chounan & Y. Yamamoto

Aza-Payne Rearrangement of Activiated 2-Aziridinemethanols and 2,3-Epoxy Amines under Basic Conditions.

J. Org. Chem., **60** (7), 2044-2058 (1995)

292. H. Tamamura, A. Otaka, J. Nakamura, K. Okubo, T. Koide, T. Ibuka & N. Fujii
Disulfide Bond-forming Reaction using a Dimethylsulfoxide/aqueous HCl
System and Its Application to Regioselective Two Disulfide Bond Formation.
Int. J. Peptide and Protein Res., **45**, 312-319 (1995)
293. H. Tamamura, T. Murakami, S. Horiuti, K. Sugihara, A. Otaka, W. Takada, T.
Ibuka, M. Waki, N. Yamamoto & N. Fujii
Synthesis of Protegrin-related Peptides and Their Antibacterial and Anti-human
Immunodeficiency Virus Activity.
Chem. Pharm. Bull., **43** (5), 853-858 (1995)
294. T. Ookura, K. Kainuma, H.-J. Kim, A. Otaka, N. Fujii & Y. Kawamura
Active Site Peptides with CXXC Motif on Map-resin can Mimic Protein Disulfide
Isomerase Activity.
Biochem. Biophys. Res. Commun., **213**(3), 746-751 (1995)
295. N. Fujii, K. Nakai, H. Tamamura, A. Otaka, N. Mimura, Y. Miwa, T. Taga, Y.
Yamamoto & T. Ibuka
SN_{2'} -Ring Opening Reaction of Aziridines Bearing an α,β-Unsaturated Ester
Group with Organocopper Reagents. A New Stereoselective Synthetic Route to
(E)-Alkene Dipeptide Isosteres.
J. Chem. Soc., Perkin I, **1995** (11), 1359-1371 (1995)
296. Otaka, K. Miyoshi, M. Kaneko, H. Tamamura, N. Fujii, M. Nomizu, T. R. Burke
Jr. & P.P. Roller
Development of Efficient Two-step Deprotection Methodology for Dimethyl-
protected Phosphoamino Acids-containing Peptide Resins and Its Application to
the Practical Synthesis of Phosphopeptides.
J. Org. Chem., **60** (13), 3967-3974 (1995)
297. K. Nakai, T. Ibuka, A. Otaka, H. Tamamura, N. Fujii & Y. Yamamoto
A One-pot Aza-Payne Rearrangement-Epoxide Ring Opening Reaction of
2-Aziridine-methanols: A Regio- and Stereoselective Synthetic Route to
Diastereomerically Pure 1,2-Amino Alcohols.
Tetrahedron Lett., **36** (35), 6247-6250 (1995)
298. Y. Kuroda, M. Ogawa, N. Fujii & T. Nakagawa
¹H-NMR Spectroscopic Studies on the Interaction between Local Anesthetics
Dibucaine and a Na⁺-Channel Inactivation Gate Peptide.
Bull. Magn. Reson., **17** (1-4), 262-263 (1995)
299. T. Kondo, H. Uenishi, T. Shimizu, T. Hirama, M. Iwashiro, K. Kurabayashi, H.
Tamamura, N. Fujii, R. Fujisawa, M. Miyazawa & H. Yamagishi

- A Single Retroviral Gag Precursor Signal Peptide Recognized by FBL-3 Tumor Specific Cytotoxic T. Lymphocytes.
J. Virol., **69** (11), 6735-6741 (1995)
300. B.S. Weeks, M. Nomizu, A. Otaka, C.A. Weston, A. Okusu, H. Tamamura & N. Fujii
The Synthetic [Tyr^{5,12}, Lys⁷]-Polyphemusin II Peptide (T22) Binds to the CD4 Cell Surface Molecule.
Biochem. Biophys. Res. Commun., **215** (2), 626-631 (1995)
301. K. Matsuzaki, K. Sugishita, N. Fujii & K. Miyajima
Molecular Basis of membrane Selectivity of an Antimicrobial Peptide, Magainin 2
Biochemistry, **34**, 3423-3429 (1995)
302. H. Tamamura, T. Murakami, M. Masuda, A. Otaka, W. Takada, T. Ibuka, H. Nakashima, M. Waki, A. Matsumoto, N. Yamamoto & N. Fujii
Structure-Activity Relationships of an Anti-HIV Peptide, T22.
Biochem. Biophys. Res. Commun., **205**, 1729-1735 (1994)
303. T. Ibuka, K. Nakai, H. Habashita, Y. Hotta, N. Fujii, N. Mimura, Y. Miwa, T. Taga & Y. Yamamoto
A Novel Route to Diastereomerically Pure (*E*)-Alkene Dipeptide Isosteres from β-Aziridinyl-α, β-enoates by Treatment with Organocopper Reagents.
Angew. Chem. Int. Ed. Engl., **33**, 652-654 (1994).
304. N. Fujii, K. Nakai, H. Habashita, Y. Hotta, H. Tamamura, A. Otaka & T. Ibuka
Synthesis of Optically Pure 2-Aziridinemethanols: Versatile Synthetic Building Blocks
Chem. Pharm. Bull., **42**, 2241-2250 (1994).
305. Otaka, H. Tamamura, Y. Terakawa, M. Masuda, T. Koide, T. Murakami, H. Nakashima, K. Matsuzaki, K. Miyajima, T. Ibuka, M. Waki, A. Matsumoto, N. Yamamoto & N. Fujii
Molecular Parameters for the Anti-Human Immunodeficiency Virus Activity of T22 ([Tyr^{5,12}, Lys⁷]-Polyphemusin II).
Biol. Pharm. Bull., **17**, 1669-1672(1994).
306. B.S. Weeks, M. Nomizu, A. Otaka, C.S. Weston, A. Okusu, H. Tamamura, A. Matsumoto, N. Yamamoto & N. Fujii
Lymphocytes and Promonocytes Attach to the Synthetic [Tyr^{5,12}, Lys⁷]-Polyphemusin II Peptide (T22).
Biochem. Biophys. Res., Commun., **202**, 470-475 (1994)
307. T. Ibuka, K. Suzuki, H. Habashita, A. Otaka, H. Tamamura, N. Mimura, Y.

- Miwa, T. Taga & N. Fujii
(*E*)-Stereoselective Synthesis of Vinylglycines from (*R*)-Serine via Organocopper-BF₃ and Related Reagents.
J. Chem. Soc., Chem. Commun., **1994**, 2151-2152 (1994)
308. K. Matsuzaki, O. Murase, H. Tokuda, S. Funakoshi, N. Fujii & K. Miyajima
Orientational and Aggregational States of Magainin 2 in Phospholipid Bilayers.
Biochemistry, **33**, 3342-3349 (1994)
309. Hosotani, R. Doi, Y. Gu, M. Wada, K. Inoue, N. Fujii, P. Rayford & M. Imamura
Metabolism of Cholecystokinin-33 in vivo: Effect of L-364718, a CCK Receptor Antagonist.
Anal. Clin. Laboratory Science, **24**, 346-354 (1994)
310. K. Sugahara, K. Shigeno, M. Masuda, N. Fujii, A. Kurosaka & K. Takeda
Structural Studies on the Chondroitinase ABC-resistant Sulfated Tetrasaccharides Isolated from Various Chondroitin Sulfate Isomers.
Carbohydr. Res., **255**, 145-163 (1994)
311. T. Shimizu, H. Uenishi, Y. Teramura, M. Iwashiro, K. Kuribayashi, H. Tamamura, N. Fujii & H. Yamagishi
Fine Structure of a Virus-Encoded Helper T-Cell Epitope Expressed on FBL-3 Tumor Cells.
J. Virol., **68**, 7704-7708 (1994)
312. Y. Kiso, N. Fujii & H. Yajima
New Disulfide Bond-Forming Reactions for Peptide and Protein Synthesis.
Brazilian J. Med. Biol. Res., **27**, 2733-274 (1994)
313. T. Ibuka, T. Taga, H. Habashita, K. Nakai, H. Tamamura, N. Fujii, Y. Chounan, H. Nemoto & Y. Yamamoto
syn-SN2' Pathway in the Reaction of Certain γ -(Mesyloxy)- α,β -Enoates with RCu(CN)MgX.BF₃ Reagents. Importance of MgX and Bulky R Group upon the Diastereoselectivity.
J. Org. Chem., **58**, 1207-1214 (1993)
314. T. Koide, H. Itoh, A. Otaka, H. Yasui, M. Kuroda, N. Esaki, K. Soda & N. Fujii
Synthetic Study on Selenocystine-Containing Peptides.
Chem. Pharm. Bull., **41**, 502-506 (1993)
315. H. Yoshizawa, A. Otaka, H. Habashita & N. Fujii
Direct Disulfide Formation from 2-Quinolinylmethyl Thioethers with Iron(III) or Copper(II) Salt.
Chem. Lett., **1993**, 803-806

316. H. Tamamura, R. Ikoma, M. Niwa, S. Funakoshi, T. Murakami & N. Fujii
Antimicrobial Activity and Conformation of Tachyplesin I and Its Analogs.
Chem. Pharm. Bull., **41**, 978-980 (1993)
317. H. Tamamura, J. Nakamura, K. Noguchi, S. Funakoshi & N. Fujii
Acceleration of *N*^ε-Deprotection Rate by the Addition of *m*-Cresol to Diluted Methanesulfonic Acid and Its Application to the Z(OMe)-Based Solid-Phase Syntheses of Human Pancreastatin-29 and Magainin 1.
Chem. Pharm. Bull., **41**, 954-957 (1993)
318. T. Koide, A. Otaka & N. Fujii
Investigation of the Dimethylsulfoxide-Trifluoroacetic acid Oxidation System for the Synthesis of Cystine-Containing Peptides.
Chem. Pharm. Bull., **41**, 1030-1034 (1993)
319. H. Tamamura, M. Kuroda, M. Masuda, A. Otaka, S. Funakoshi, H. Nakashima, N. Yamamoto, M. Waki, A. Matsumoto, J.M. Lancelin, D. Kohda, S. Tate, F. Inagaki & N. Fujii
A Comparative Study of the Solution Structures of Tachyplesin I and a Novel Anti-HIV Synthetic Peptides, T22([Tyr_{5,12}, Lys₇]-Polyphemusin II), Determined by Nuclear Magnetic Resonance.
Biochim. Biophys. Acta., **1163**, 209-216 (1993)
320. H. Tamamura, A. Otaka, J. Nakamura, K. Okubo, T. Koide, K. Ikeda & N. Fujii
Disulfide Bond Formation in S-Acetamidomethyl Cysteine-Containing Peptides by the Combination of Silver Trifluoromethanesulfonate and Dimethylsulfoxide/Aqueous HCl
Tetrahedron Lett., **34**, (31), 4931-4934 (1993)
321. S. Funakoshi, H. Fukuda & N. Fujii
Affinity Purification Method using a Reversible Biotinylation Reagent for Peptides Synthesized by the Solid-Phase Technique.
J. Chromatography, **638**, 21-27 (1993)
322. N. Fujii, K. Nakai, H. Habashita, H. Yoshizawa, T. Ibuka, F. Garrido, A. Mann, Y. Chounan, & Y. Yamamoto
SN 2' Selective Alkylation of Allylic Chlorides and Mesylates with RZnX Reagents Generated from Grignard Reagents, Zinc Chloride, Lithium Chloride and Cu(II)-Salts.
Tetrahedron Lett., **34**, (26), 4227-4230 (1993)
323. M. Iwashiro, T. Kondo, T. Shimizu, H. Yamagishi, K. Takahashi, Y. Matsubayashi, T. Masuda, A. Otaka, N. Fujii, A. Ishoto, M. Miyazawa, M.N.

- Robertson, B. Chesebro & K. Kuribayashi
Multiplicity of Virus-Encoded Helper T-Cell Epitopes Expressed on FBL-3
Tumor Cells.
J. Virology, **67** (8), 4533-4542 (1993).
324. T. Koide, H. Itoh, A. Otaka, M. Furuya, Y. Kitajima & N. Fujii
Syntheses and Biological Activities of Selenium Analogs of α -Rat Atrial
Natriuretic Peptide.
Chem. Pharm. Bull., **41** (9), 1596-1600 (1993)
325. Y. Kasuya, K. Fujimoto, M. Miyamoto, T. Juji, A. Otaka, S. Funakoshi, N. Fujii
& H. Kawaguchi
Preparation of Peptide-carrying Microspheres with Bioactivity on Platelets.
J. Biomater. Sci. Polymer Edn., **4** (4), 369-380 (1993).
326. K. Matsuzaki, M. Fukui, N. Fujii & K. Miyajima
Permeabilization and Morphological Changes in Phosphatidylglycerol Bilayers
Induced by an Antimicrobial Peptide, Tachyplesin I.
Colloid & Polymer Sciences, **271**, 901-908 (1993).
327. T. Ibuka, K. Nakai, H. Habashita, N. Fujii, F. Garrido, A. Mann, Y. Chounan &
Y. Yamamoto
Unprecedented Rearrangement Reaction of 2-Aziridinemethanols with "Lower
Order" Lithium Methylcyanocuprate
Tetrahedron Lett., **34** (46), 7421-7424 (1993)
328. T. Ibuka, K. Nakai, H. Habashita, K. Bessho, N. Fujii, Y. Chounan & Y.
Yamamoto,
Remarkable Differences in Reactivity of Ordinary Vinylcopper Reagents and
Vinylzinc Halide Containing a Copper Salt towards γ -Mesyloxy- α,β -Enoates.
Synthesis of Homochiral 1,4-Dienes
Tetrahedron Lett., **49** (42) 9479-9488 (1993).
329. K. Matsuzaki, M. Nakayama, M. Fukui, A. Otaka, S. Funakoshi, N. Fujii, K.
Bessho & K. Miyajima
Role of Disulfide Linkage in Tachyplesin-Lipid Interaction.
Biochemistry, **32** (43), 11704-11710 (1993).
330. H. Nakashima, Y. Yamamoto, M. Masuda & N. Fujii
Defensins Inhibit HIV Replication in vitro.
AIDS, **7**, 1129 (1993)
331. T. Tamura, T. Oikawa, A. Otaka, N. Fujii, N. Esaki & K. Soda
Synthesis and Characterization of the Selenium Analog of Glutathione Disulfide

- Analytical Biochemistry*, **202**, 151-154 (1993).
332. Funakoshi, K. Miyasaka, K. Kitani, J. Nakamura, S. Funakoshi, H. Fukuda & N. Fujii
Stimulatory Effects on Islet Amyloid Polypeptide (Amylin) on Exocrine Pancreas and Gastrin Release in Conscious Rats.
Regul. Peptides., **38**, 135-143 (1992)
333. Kohasi, T. Ono, T. Soejima, T. Moriya, A. Umeda, Y. Meno, K. Amako, S. Funakoshi, M. Masuda & N. Fujii
Bactericidal Activities of Rat Defensins and Synthetic Rabbit Defensins on *Staphylococci*, *Klebsiella Pneumoniae* (Chedid, 277, and 8N3), *Pseudomonas Aeruginosa* (Mucoid and Nonmucoid Strains), *Salmonella Typhimurium* (Ra, Rc, Rd, and Re of LPS Mutants) and *Escherichia Coli*.
Microbiol. Immunol., **36**, 369-380 (1992)
334. Yonezawa, J. Kuwahara, N. Fujii & Y. Sugiura
Binding of Tachyplesin I to DNA Revealed by Footprinting Analysis. Significant Contribution of Secondary Structure to DNA Binding and Implication for Biological Action.
Biochemistry, **31**, 2998-3004 (1992)
335. H. Nakashima, M. Masuda, T. Murakami, Y. Koyanagi, A. Matsumoto, N. Fujii & N. Yamamoto
Anti-Human Immunodeficiency Virus Activity of a Novel Synthetic Peptide, T22 ([Tyr-5,12, Lys7]Polypheusin II): a Possible Inhibitor of Virus-Cell Fusion.
Antimicrobial Agents and Chemotherapy, **36**, 1249-1255 (1992)
336. T. Ibuka, H. Yoshizawa, H. Habashita, N. Fujii, Y. Chounan, M. Tanaka & Y. Yamamoto
"Higher Order" Zinc Cuprates Involving Lithium Chloride: Synthesis of (*E*)-Alkene Dipeptide Isosteres Free From Reductive Elimination Products.
Tetrahedron Lett., **26**, 3783-3786 (1992)
337. M. Masuda, H. Nakashima, T. Ueda, H. Naba, R. Ikoma, S. Funakoshi, T. Murakami, Y. Koyanagi, N. Yamamoto & N. Fujii
Structure-Activity Relationships of Tachyplesin Analogs as Anti-HIV Agents.
J. Pharmacobio-Dyn., **15**, s-90 (1992)
338. M. Masuda, H. Nakashima, T. Ueda, H. Naba, R. Ikoma, A. Otaka, Y. Terakawa, H. Tamamura, T. Ibuka, T. Murakami, Y. Koyanagi, M. Waki, A. Matsumoto, N. Yamamoto, S. Funakoshi & N. Fujii
A Novel Anti-HIV Synthetic Peptide, T-22 ([Tyr5,12, Lys7]- Polypheusin II).

Biochem. Biophys. Res. Commun., **189**, 845-850 (1992)

339. N. Fujii, C.A.S.A. Minetti, H. L. Nakhasi, S.W. Chen, E. Barbehenn, P.H. Nunes, and N. Y. Nguyen
Isolation, cDNA Cloning, and Characterization of an 18-kDa Hemagglutinin and Amebocyte Aggregation Factor from *Limulus Polyphemus*.
J. Biol. Chem., **267**, 22452-22459 (1992).
340. Y. Kuraishi, M. Kawamura, T. Yamaguchi, T. Houtani, S. Kuwabata, S. Futaki, N. Fujii & M. Satoh
Intrathecal Injection of Galanin and Its Antiserum Affect nociceptive Response of Rat to Mechanical, but not Thermal, Stimuli.
Pain., **44**, 321-324 (1991)
341. K. Matsuzaki, M. Harada, S. Funakoshi, N. Fujii & K. Miyajima
Physicochemical Determinants for the Interaction of Magainin 1 and 2 with Acidic Lipid Bilayers.
Biochim. Biophys. Acta., **1063**, 162-170 (1991)
342. S. Funakoshi, H. Fukuda & N. Fujii
Chemoselective One-Step Purification Method for Peptides Synthesized by the Solid-Phase Technique.
Proc. Natl. Acad. Sci. USA., **88**, 6981-6985 (1991)
343. T. Ibuka, H. Habasha, A. Otaka, N. Fujii, Y. Oguchi, T. Uyehara & Y. Yamamoto
A Highly Stereoselective Synthesis of (*E*)-Alkene Dipeptide Isosteres via Organocyanato-copper-Lewis Acid Mediated Reaction.
J. Org. Chem., **56**, 4370-4382 (1991)
344. N. Fujii, H. Habashita, N. Shigemori, A. Otaka, T. Ibuka, M. Tanaka & Y. Yamamoto
A Simple Synthesis of $\psi[(E)\text{CH}=\text{CH}]$ Gly Dipeptide Isosteres via Reductive Elimination of γ -Oxygenated α,β -Enoates with Alkenylcopper Reagents.
Tetrahedron Lett., **32**, 4969-4972 (1991)
345. Otaka, T. Koide, A. Shide & N. Fujii
Application of Dimethylsulphoxide(DMSO)/Trifluoroacetic Acid (TFA) Oxidation to the Synthesis of Cystine-Containing Peptide.
Tetrahedron Lett., **32**, 1223-1226 (1991)
346. T. Koide, A. Otaka, H. Suzuki & N. Fujii
Selective Conversion of *S*-Protected Cysteine Derivatives to Cystine by Various Sulphoxide-Silyl Compound / Trifluoroacetic Acid Systems.
Synlett., **1991**, 345-346

347. M. Nomizu, Y. Inagaki, T. Yamashita, A. Ohkubo, A. Otaka, N. Fujii, P.P. Roller & H. Yajima
Two-Step Hard Acid Deprotection / Cleavage Procedure for Solid Phase Peptide Synthesis.
Int. J. Peptide Protein Res., **37**, 145-152(1991)
348. M. Nomizu, Y. Inagaki, A. Iwamatsu, T. Kashiwabara, H. Ohta, A. Morita, K. Nishikori, A. Otaka, N. Fujii & P.P. Roller
Solid Phase Peptide Synthesis of Human Endothelin Precursor Peptides Using Two-Step Hard Acid Deprotection/Cleavage Methods.
Int. J. Peptide Protein Res., **38**, 580-587 (1991)
349. K. Matsuzaki, M. Fukui, N. Fujii & K. Miyajima
Interaction of an Antimicrobial Peptide, Tachyplesin I, with Lipid Membranes.
Biochim. Biophys. Acta., **1070**, 259-264 (1991)
350. H. Shinozaki, K. Miyasaka, H. Wakasugi, N. Fujii & A. Funakoshi
Bioactivity of synthetic human cholecystokinin (CCK)-33 in vitro and in vivo.
Gastroenterol. Jpn., **26**, 51-55 (1991)
351. K. Miyasaka, A. Funakoshi, K. Kitani, H. Tamamura, N. Fujii & S. Funakoshi
The Importance of the CTerminal Amide Structure of Rat Pancreastatin to Inhibit Pancreatic Exocrine Secretion.
FEBS Lett., **263**, 279-280 (1990)
352. H. Tamamura, M. Ohta, K. Yoshizawa, Y. Ono, A. Funakoshi, K. Miyasaka, K. Tateishi, A. Jimi, H. Yajima, N. Fujii & S. Funakoshi
Isolation and Characterization of a Tumor-Derived Human Protein Related to Chromogranin A and Its in Vitro Conversion to Human Pancreastatin-48.
Eur. J. Biochem., **191**, 33-39 (1990)
353. K. Miyasaka, A. Funakoshi, K. Kitani, H. Tamamura, S. Funakoshi & N. Fujii
Inhibitory Effect of Pancreastatin on Pancreatic Exocrine Secretion. Pancreastatin Inhibits Central Vagal Nerve Stimulation.
Gastroenterology, **99**, 1751-1756 (1990)
354. K. Miyasaka, A. Funakoshi, Y. Yasunami, R. Nakamura, K. Kitani, H. Tamamura, S. Funakoshi & N. Fujii
Rat Pancreastatin Inhibits Both Pancreatic Exocrine and Endocrine Secretions in Rats.
Regulatory Peptides, **28**, 189-198 (1990)
355. M. Takeyama, N. Yanaga, K. Yarimizu, J. Ono, R. Takaki, N. Fujii & H. Yajima
Enzyme Immunoassay of Somatostatin (SS)-like Immunoreactive Substance in

- Bovine Milk.
- Chem. Pharm. Bull.*, **38**, 456-459 (1990)
356. M. Takeyama, K. Wakayama, F. Takayama, K. Kondo, N. Fujii & H. Yajima
Micro-Enzyme Immunoassay of Vasoactive Intestinal Polypeptide (VIP)-like
Immunoreactive Substance in Bovine Milk.
Chem. Pharm. Bull., **38**, 960-962 (1990)
357. M. Takeyama, F. Yasunaga, A. Otaka, N. Fujii & H. Yajima
Microenzyme Immunoassay for the Measurement of Brain Natriuretic Peptide
(BNP)-Like Immunoreactivity in Porcine Plasma.
J. Immunol. Methods., **130**, 217-222 (1990)
358. T. Tominaga, J. Fukata, Y. Naito, Y. Nakai, S. Funakoshi, N. Fujii, & H. Imura
Effects of Corticostatin-I on Rat Adrenal Cells in Vitro.
J. Endocrinol., **125**, 287-292 (1990)
359. T. Ibuka, H. Habashita, S. Funakoshi, N. Fujii, K. Baba, M. Kozawa, Y. Oguchi,
T. Uyehara & Y. Yamamoto
Determination of Absolute Configuration of the Alkyl Group at the α -Position in
the Acyclic α -Alkyl-(*E*)- β,γ -Enoates by Circular Dichroism.
Tetrahedron Assym., **1**, 389-394 (1990)
360. T. Ibuka, H. Habashita, S. Funakoshi, N. Fujii, Y. Oguchi, T. Uyehara & Y.
Yamamoto
Highly Selective Synthesis of (*E*)-Alkene Isosteric Dipeptides with High Optical
Purity via $\text{RCu}(\text{CN})\text{Li} \cdot \text{BF}_3$ Mediated Reaction.
Angew. Chem. Int. Ed. Engl., **29**, 801- 803 (1990)
361. Y. Yoshihara, H. Ueda, N. Fujii, A. Shide, H. Yajima & M. Satoh
Purification of a Novel Type of Calcium-activated Neutral Protease from Rat
Brain .
J. Biol. Chem., **265** (10), 5809-5815 (1990)
362. M. Takeyama, K. Mori, F. Takayama, K. Kondo, K. Kitagawa & N. Fujii
Enzume Immunoassay of a Substance P-like Immunoreactive Substance in
Human Plasma and Saliva.
Chem. Pharm. Bull., **38** (12), 3494-3496 (1990)
363. R. Doi, R. Hosotani, K. Inoue, M. Kogire, S. Sumi, N. Fujii, H. Yajima, P.L.
Rayford & T. Tobe
Effect of Synthetic Human CCK-33 on Pancreatic Blood Flow in Dogs.
Pancreas, **5**, 615-620 (1990)
364. R. Doi, R. Hosotani, K. Inoue, N. Fujii, H. Yajima, P.L. Rayford & T. Tobe

- Receptor Binding of Cholecystokinin Analogues in Isolated Rat Pancreatic Acini.
Biochem. Biophys. Res. Commun., **166**, 286-292 (1990)
365. K. Sugahara, M. Nakamura, J. Nagisa, M. masuda, Y. Nunokawa, N. Fujii & I. Yamashina
Regulation of Serum Glycosaminoglycan Sulfotransferase Activities; Inhibition by sulfated Glycosaminoglycans and Activation by Polyamines and Basic Peptides Including a Polylysine-containing Segment of the c-Ki-ras 2 Protein.
J. Biochem., **106**, 910-919 (1989)
366. K. Takaori, K. Inoue, M. Kogire, R. Doi, S. Sumi, M. Yun, N. Fujii, H. Yajima & T. Tobe
Effect of Synthetic Physalaemin on Splanchnic Circulation in Dogs.
Life Science, **44**, 667-672 (1989)
367. Funakoshi, K. Miyasaka, K. Kitani, S. Funakoshi, H. Tamamura, N. Fujii, I. Nakano & K. Tatemoto
Comparative Effects of Mamalian Pancreastatins on the Pancreatic Exocrine Secretion.
Jpn. J. Physiol., **39**, 901-905 (1989)
368. H. Yamada, M. Tsutsumi, M. Fukase, A. Fujimori, Y. Yamamoto, A. Miyauchi, Y. Fujii, T. Noda, N. Fujii & T. Fujita
Effects of Human PTH-Related Peptide and Human PTH on Cyclic-AMP Production and Cytosolic Free Calcium in an Osteoblastic Cell Clone.
Bone and Mineral., **6**, 45-54 (1989)
369. Y. Kuraishi, T. Nakayama, H. Ohno, N. Fujii, A. Otaka, H. Yajima & M. Satoh
Calcitonin Gene-Related Peptide Increases in the Dorsal Root Ganglia of Adjuvant Arthritic Rat.
Peptides., **10**, 447-452(1989)
370. M. Takeyama, M. Hatano, A. Otaka, N. Fujii & H. Yajima
Immuno-affinity Purification of Specific Antibodies against Vasoactive Intestinal Polypeptide (VIP) by the VIP(1-10)-linked Polydimethylacrylamide Resin.
Chem. Pharm. Bull., **37**, 834-837(1989)
371. Otaka, H. Morimoto, N. Fujii, T. Koide, S. Funakoshi & H. Yajima
S-Benzylloxymethylcysteine, Its Properties and Application in the Synthesis of Porcine Brain Natriuretic Peptide (pBNP).
Chem. Pharm. Bull., **37**, 526-528 (1989)
372. K. Matsuzaki, M. Harada, T. Handa, S. Funakoshi, N. Fujii, H. Yajima & K. Miyajima

- Magainin 1-Induced Leakage of Entrapped Calcein Out of Negatively-Charged Lipid Vesicles.
Biochim. Biophys. Acta, 981, 130-134 (1989)
373. Y. Hayashi, K. Ikuta, N. Fujii, K. Ezawa & S. Kato
Inhibition of HIV-1 Replication and Syncytium Formation by Synthetic CD-4 Peptides.
Arch. Virol., **105**, 129-135 (1989)
374. K. Akaji, Y. Hayashi, N. Fujii, T.Y. Liu, I. Berkower & H. Yajima
Solid-Phase Syntheses of Immunological Properties of Fragment Peptides Related to Human Malaria Circumsporozoite Protein.
Chem. Pharm. Bull., **37**, 1612-1615 (1989)
375. M. Takeyama, T. Kino, L. Guo, A. Otaka, N. Fujii & H. Yajima
Immunoaffinity Purification of Specific Antibodies Against Human Gastrin Releasing Peptide (h-GRP) by the h-GRP(1-8)-linked Polydimethylacrylamide Resin.
Int. J. Peptide Protein Res., **33**, 457-462 (1989)
376. N. Fujii, A. Otaka, T. Watanabe, A. Okamachi, H. Tamamura, H. Yajima, Y. Inagaki, M. Nomizu & K. Asano
Silver Trifluoromethanesulphonate as an *S*-Deprotecting Reagent for the Synthesis of Cystine Peptides.
J. Chem. Soc., Chem. Commun., **1989**, 283-284
377. K. Akaji, N. Fujii, F. Tokunaga, T. Miyata, S. Iwanaga & H. Yajima
Syntheses of Three Peptides Isolated from Horseshoe Crab Hemocytes, Tachyplesin I, Tachyplesin II, and PolypheMusin I.
Chem. Pharm. Bull., **37**, 2661-2664 (1989)
378. S. Funakoshi, H. Tamamura, M. Ohta, K. Yoshizawa, A. Funakoshi, K. Miyasaka, K. Tateishi, K. Tatemoto, I. Nakano, H. Yajima & N. Fujii
Isolation and Characterization of a Tumor-Derived Human Pancreastatin-Related Protein.
Biochem. Biophys. Res. Commun., **164**, 141-148 (1989)
379. R. Doi, K. Inoue, M. Kogure, S. Sumi, M. Yun, S. Futaki, N. Fujii, H. Yajima & T. Tobe
Effect of Synthetic Human Cholecystokinin-33 on Exocrine Pancreas.
Biochem. Biophys. Res. Commun., **150**, 1251-1255 (1988)
380. N. Fujii, S. Futaki, H. Morimoto, K. Inoue, R. Doi, T. Tobe & H. Yajima
Total Synthesis of Human Cholecystokinin-33.

J. Chem. Soc., Chem. Commun., **1988**, 324-325

381. N. Fujii, S. Futaki, S. Funakoshi, K. Akaji, H. Morimoto, O. Ikemura & H. Yajima
Preparation of a Protected 33-Residue Peptide for the Synthesis of Human Cholecystokinin (hCCK-33).
Chem. Pharm. Bull., **36**, 3271- 3280 (1988).
382. N. Fujii, S. Futaki, S. Funakoshi, K. Akaji, H. Morimoto, R. Doi, K. Inoue, M. Kogire, S. Sumi, M. Yun, T. Tobe, M. Aono, M. Matsuda, H. Narusawa, M. Moriga & H. Yajima
Synthesis of a 33-Residue Peptide Corresponding to the Entire Amino Acid Sequence of Human Cholecystokinin (hCCK-33).
Chem. Pharm. Bull., **36**, 3281-3291 (1988)
383. H. Yajima, N. Fujii, S. Funakoshi, T. Watanabe, E. Murayama & A. Otaka
New Strategy for the Chemical Synthesis of Proteins.
Tetrahedron, **44**, 805-819 (1988).
384. N. Fujii, T. Watanabe, T. Aotake, A. Otaka, I. Yamamoto, J. Konishi & H. Yajima
Synthesis of Chicken Calcitonin-Gene-Related Peptide (cCGRP) by Application of Sulfoxide-Directed Disulfide-Bond Forming Reaction.
Chem. Pharm. Bull., **36**, 3304-3311 (1988)
385. M. Hatano, S. Funakoshi, N. Fujii, M. Takeyama, M. Yun, K. Inoue, M. Kogire, T. Tobe & H. Yajima
Synthesis of Guinea Pig Vasoactive Intestinal Polypeptide (gVIP).
Chem. Pharm. Bull., **36**, 3857-3866 (1988)
386. S. Funakoshi, E. Murayama, L. Guo, N. Fujii & H. Yajima
A Modified Benzhydrylamine as a Handle Reagent for the Solid Phase Synthesis of Peptide Amide Based on the Fluorenylmethoxycarbonyl Method.
J. Chem. Soc., Chem. Commun., **1988**, 382-384
387. S. Funakoshi, H. Tamamura, N. Fujii, K. Yoshizawa, H. Yajima, K. Miyasaka, A. Funakoshi, M. Ohta, Y. Inagaki & L.A. Carpino
Combination of a New Amide-Precursor Reagent and Trimethylsilyl Bromide Deprotection for the Fmoc-based Solid Phase Synthesis of Human Pancreastatin and One of Its Fragments.
J. Chem. Soc., Chem. Commun., **1988**, 1588-1590
388. A. Funakoshi, K. Miyasaka, R. Nakamura, K. Kitani, S. Funakoshi, H. Tamamura, N. Fujii & H. Yajima

- Bioactivity of Synthetic Human Pancreastatin on Exocrine Pancreas.
Biochem. Biophys. Res. Commun., **156**, 1237-1242 (1988)
389. L. Guo, S. Funakoshi, N. Fujii & H. Yajima
Combination of a New Amide-precursor Reagent and Trimethylsilylchloride
Deprotection for the 9-Fluorenylmethyloxycarbonyl-Based Solid Phase
Synthesis of Chicken Antral Peptide.
Chem. Pharm. Bull., **36**, 4989-4992 (1988)
390. L. Guo, E. Murayama, S. Funakoshi, N. Fujii, M. Aono, M. Matsuda, M. Moriga
& H. Yajima
Solution Phase Synthesis of a 36-Residue Peptide Amide Corresponding to the
Entire Amino Acid Sequence of Chicken Antral Peptide.
Chem. Pharm. Bull., **36**, 4364-4376 (1988)
391. C. Shigeno, I. Yamamoto, N. Kitamura, T. Noda, K. Lee, T. Sone, K. Shiomi, A.
Otaka, N. Fujii, H. Yajima & J. Konishi
Interaction of Human Parathyroid Hormone-related Peptide with Parathyroid
Hormone Receptors in Clonal Rat Osteosarcoma Cells.
J. Biol. Chem., **263**, 18369-18377 (1988)
392. Y. Hayashi, S. Katakura, M. Nomizu, A. Tashiro, S. Kuwata, N. Miaki, T.Y. Liu,
N. Fujii & H. Yajima
Solid-Phase Syntheses and Immunological Properties of Fragment Peptides
Related to Human Hepatitis B Virus Surface Antigen (HBsAg) and Its
Pre-S2 Gene.
Chem. Pharm. Bull., **36**, 4993-4999 (1988)
393. K. Sakina, N. Fujii, S. Funakoshi & H. Yajima
Partial Structural Revision of Porcine Pancreatic Colipase.
Chem. Pharm. Bull., **36**, 424-429 (1988)
394. N. Fujii, A. Otaka, S. Funakoshi, T. Watanabe, H. Arai, K. Bessho & H. Yajima
Model Experiments for the Synthesis of Open-Chain Unsymmetrical Cystine
Peptides.
J. Protein Chemistry., **7**, 151-156 (1988)
395. S. Funakoshi, E. Murayama, L. Guo, N. Fujii & H. Yajima
A Modified Benzhydrylamine, as a Useful Handle Reagent for the Fmoc-Based
Solid Phase Synthesis of Peptides Amides.
Coll. Czech. Chem. Commun., **53**, 2791-2800 (1988)
396. M. Nakai, M. Fukase, K. Sakaguchi, T. Noda, N. Fujii & T. Fujita
Human Parathyroid Hormone Related Protein Fragment-(1-34) Had

- Glucose-6-Phosphate Dehydrogenase Activity of Distal Convolved Tubules in Cytochemical Bioassay.
Biochem. Biophys. Res. Commun., **154**, 146-150 (1988)
397. Fujimori, M. Tsutsumi, H. Yamada, M. Fukase, T. Noda, N. Fujii & T. Fujita
The Human Parathyroid Hormone-Related Protein Mimics the Action of Human Parathyroid Hormone in an Osteoblastic Cell Line, MC3T3-E1.
JBMM(日本骨代謝学会雑誌), **6**, 172-175 (1988)
398. M. Kogire, K. Inoue, S. Sumi, R. Doi, K. Takaori, M. Yun, N. Fujii, H. Yajima & T. Tobe
Effects of Synthetic Human Gastric Inhibitory Polypeptide on Splanchnic Circulation in Dogs.
Gastroenterology, **95**, 1636-1640 (1988)
399. N. Fujii, A. Otaka, O. Ikemura, K. Akaji, S. Funakoshi, Y. Hayashi, Y. Kuroda & H. Yajima
Trimethylsilyl Trifluoromethanesulfonate as a Useful Deprotecting Reagent in Both Solution and Solid Phase Synthesis.
J. Chem. Soc. Chem. Commun., **1987**, 274-275 (1987)
400. N. Fujii, A. Otaka, O. Ikemura, M. Hatano, A. Okamachi, S. Funakoshi, M. Sakurai, T. Shioiri & H. Yajima
Hard Acid Deprotecting Procedure for Peptide Synthesis.
Chem. Pharm. Bull., **35**, 3447-3452 (1987)
401. N. Fujii, A. Otaka, N. Sugiyama, M. Hatano & H. Yajima
Evaluation of Trimethylsilyl Bromide as a Hard Acid Deprotecting Reagent in Peptide Synthesis.
Chem. Pharm. Bull., **35**, 3880-3883 (1987)
402. F. A. Robey, K. Ohura, S. Futaki, N. Fujii, H. Yajima, N. Goldman, K.D. Jones & S. Wahl
Proteolysis of Human C-Reactive Protein Produces Peptides with Potent Immuno-modulating Activity.
J. Biol. Chem., **262**, 7053-7057 (1987)
403. N. Fujii, A. Otaka, S. Funakoshi, K. Bessho & H. Yajima
New Procedure for the Synthesis of Cystine-Peptides by Oxidation of *S*-Substituted Cysteine-Peptides with Thallium(III) Trifluoroacetate.
J. Chem. Soc., Chem. Commun., **1987**, 163-164 (1987)
404. N. Fujii, A. Otaka, S. Funakoshi, K. Bessho, T. Watanabe, K. Akaji & H. Yajima
Synthesis of Cystine Peptides by Oxidation of *S*-Protected Cysteine Peptides

- with Thallium (III) Trifluoroacetate.
Chem. Pharm. Bull., **35**, 2339-2347 (1987)
405. N. Fujii, A. Otaka, T. Watanabe, H. Arai, S. Funakoshi, K. Bessho & H. Yajima Sulphoxide-Directed Disulphide Bond-Forming Reaction for the Cystine Peptides.
J. Chem. Soc., Chem. Commun., **1987**, 1676-1678 (1987)
406. N. Fujii, Y. Hayashi, K. Akaji, S. Funakoshi, M. Shimokura, S. Yugushi, L. H. Lazarus & H. Yajima Solid Phase Synthesis of a Rabbit Stomach Peptide by Application of a New Polymer Support and a New Deprotecting Procedure.
Chem. Pharm. Bull., **35**, 1266-1269 (1987)
407. Akaji, M. Nomizu, K. Watanabe, S. Funakoshi, H. Imura, T. Tsukada, J. Fukata, N. Nakamura, M. Koida, N. Fujii & H. Yajima Application of the Hard Acid Deprotecting Procedure to the Synthesis of Ovine Corticotropin Releasing Factor (oCRF).
Chem. Pharm. Bull., **35**, 3859-3865 (1987)
408. Fujii, T. Watanabe, A. Otaka, K. Bessho, I. Yamamoto, T. Noda & H. Yajima Synthesis of Second Human Calcitonin Gene-Related Peptide (β -hCGRP) by Application of a New Disulfide-Bonding Reaction with Thallium (III) Trifluoroacetate.
Chem. Pharm. Bull., **35**, 4769-4776 (1987)
409. R. Oku, M. Satoh, N. Fujii, A. Otaka, H. Yajima & H. Takagi Calcitonin Gene-Related Peptide Promotes Mechanical Nociceptin by Potentiating Release of Substance P from the Spinal Dorsal Horn in Rats.
Brain Research., **403**, 350-354 (1987)
410. Fujii, O. Ikemura, S. Funakoshi, H. Matsuo, T. Segawa, Y. Nakata, A. Inoue & H. Yajima Application of a New Deprotecting Procedure with Trimethylsilyl Trifluoromethane-sulfonate for the Synthesis of Two Porcine Spinal Cord Peptides, Neuromedin U-8 and Neuromedin U-25.
Chem. Pharm. Bull., **35**, 1076-1084 (1987)
411. Sugiyama, N. Fujii, S. Funakoshi, A. Funakoshi, K. Miyasaka, M. Aono, M. Moriga, K. Inoue, M. Kogure, S. Sumi, R. Doi, T. Tobe & H. Yajima Synthesis of a 36-Residue Peptide Amide Corresponding to the Entire Amino Acid Sequence of Human Pancreatic Polypeptide.
Chem. Pharm. Bull., **35**, 3585-3596 (1987)

412. M.F. Good, J.A. Berzofsky, W.L. Malov, Y. Hayashi, N. Fujii, W.T. Hockmeyer & L.A. Miller
Genetic Control of the Immune Response in Mice to a *Plasmodium Faliparum* Sporozoite Vaccine; Widespread Nonresponsiveness to Single Malaria T Epitope in Highly Repetitive Vaccine.
J. Exp. Med., **164**, 655-660 (1986)
413. H. Yajima, S. Futaki, N. Fujii, K. Akaji, S. Funakoshi, M. Sakurai, S. Katakura, K. Inoue, R. Hosotani, T. Tobe, T. Segawa, A. Inoue, K. Tatemoto & V. Mutt
Synthesis and Biological Activity of Galanin, a Novel Porcine Intestinal Polypeptide.
Chem. Pharm. Bull., **34**, 528-539 (1986).
414. S. Kuno, W. Li, N. Fujii, Y. Kai, H. Adachi, K. Bessho, T. Segawa, Y. Nakata, T. Inoue & H. Yajima
Synthesis of Gln15-Motilin and Examination of Its Immunological Properties.
Chem. Pharm. Bull., **34**, 4811-4816 (1986)
415. N. Fujii, M. Sakurai, K. Akaji, M. Nomizu, H. Yajima, K. Mizuta, M. Aono, M. Moriga, K. Inoue, R. Hosotani & T. Tobe
Solution Synthesis of a 42-Residue Peptide Corresponding to the Entire Amino Acid Sequence of Human Glucose-Dependent Insulinotropic Polypeptide (GIP).
Chem. Pharm. Bull., **34**, 2397-2410 (1986)
416. M. Sakurai, K. Akaji, N. Fujii, M. Moriga, M. Aono, K. Mizuta, H. Adachi, M. Noguchi, K. Inoue, R. Hosotani, T. Tobe & H. Yajima
Synthesis of a 42-Residue Peptide Corresponding to the Entire Amino Acid Sequence of Porcine GIP (Glucose-Dependent Insulinotropic Polypeptide).
Chem. Pharm. Bull., **34**, 3447-3453 (1986)
417. N. Fujii, A. Otaka, S. Funakoshi, M. Nomizu, K. Akaji, H. Yajima, K. Kitagawa, T. Akita, K. Ando & T. Kawamoto
Preparation of Seven Peptide Fragments for the Synthesis of Human Calcitonin Gene-Related Peptide (hCGRP).
Chem. Pharm. Bull., **34**, 603-612 (1986)
418. N. Fujii, A. Otaka, S. Funakoshi, M. Nomizu, K. Akaji, H. Yajima, I. Yamamoto, K. Torizuka, K. Kitagawa, T. Akita, K. Ando, T. Kawamoto, Y. Shimonishi & T. Takao
Solution Phase Synthesis of a 37-Residue Peptide Amide Corresponding to the Entire Amino Acid Sequence of Human Calcitonin Gene-Related Peptide (hCGRP).

- Chem. Pharm. Bull.*, **34**, 613-620 (1986)
419. Otaka, N. Fujii, S. Funakoshi, I. Yamamoto, K. Torizuka & T. Noda
Synthesis of Des-1-Ala-des- α -Amino-Human Calcitonin Gene-Related Peptide.
Chem. Pharm. Bull., **34**, 3915-3918 (1986)
420. N. Fujii, M. Nomizu, S. Futaki, A. Otaka, S. Funakoshi, K. Akaji, K. Watanabe & H. Yajima
Evaluation of Two- β -Carboxyl Protecting Group of Aspartic Acid, Cyclohexyl and Cyclooctyl, for Peptide Synthesis.
Chem. Pharm. Bull., **34**, 864-868 (1986)
421. H. Yajima, S. Futaki, A. Otaka, T. Yamashita, S. Funakoshi, K. Bessho, N. Fujii & K. Akaji
Evaluation of β -Menthylaspartate for Peptide Synthesis.
Chem. Pharm. Bull., **34**, 4356-4361 (1986)
422. M. Shimokura, Y. Kiso, A. Nagata, M. Tsuda, H. Seki, Y. Kai, N. Fujii & H. Yajima
Conventional Solution Phase Synthesis of Porcine Hypothalamic Growth Hormone Releasing Factor (pGRF).
Chem. Pharm. Bull., **34**, 1814-1820 (1986)
423. M. Shimokura, Y. Kiso, A. Nagata, M. Tsuda, H. Seki, Y. Kai & N. Fujii
Conventional Solution Synthesis of Bovine Hypothalamic Growth Hormone Releasing Factor (bGRF).
Chem. Pharm. Bull., **34**, 2218-2228 (1986)
424. H. Nakamura, Y. Fukuda, M. Koide, N. Fujii, A. Otaka, S. Funakoshi, H. Yajima, N. Mitsuyasu & R.C. Orlowski
Binding Site of Calcitonin Gene-Related Peptide (CGRP): Abundant Occurrence in Visceral Organs.
Japan J. Pharmacol., **42**, 175-180 (1986)
425. I. Yamamoto, N. Kitamura, J. Aoki, C. Shigeno, M. Hino, K. Asonuma, K. Torizuka, N. Fujii, A. Otaka & H. Yajima
Human Calcitonin Gene-related Peptide Possesses Weak Inhibitory Potency of Bone Resorption *in vitro*.
Calcif. Tissue Int., **38**, 339-341 (1986)
426. M. Satoh, R. Oku, A. Maeda, N. Fujii, A. Otaka, S. Funakoshi, H. Yajima & H. Takagi
Possible Mechanism of Positive Inotropic Action of Synthetic Human Calcitonin Gene-Related Peptide in Isolated Rat Atrium.

Peptides, **7**, 631-635 (1986)

427. N. Fujii, S. Kuno, A. Otaka, S. Funakoshi, K.Takagi & H. Yajima
Conventional Procedure for the Reduction of Methionine Sulfoxide.
Chem. Pharm. Bull., **33**, 4587-4588 (1985)
428. N. Fujii, M. Sakurai, S. Kuno, H. Yajima, M. Satoh, M. Matsushita, N. Yamamoto, H. Takagi, Z.M. Wang, W. Lee & P.F. Wang
Synthesis of Adrenorphin and Enkephalin Analogs.
Chem. Pharm. Bull., **33**, 4326-4332 (1985)
429. N. Fujii, H. Yajima, A. Otaka, S. Funakoshi, M. Nomizu, K. Akaji, I. Yamamoto, K. Torizuka, K. Kitagawa, T. Akita, K. Ando, T. Kawamoto, Y. Shimonishi & T.Takao
Application of *S*-1-Adamantylcysteine to the Synthesis of a 37-residue Peptide Amide Corresponding to Human Calcitonin Gene-Related Peptide (hCGRP).
J. Chem. Soc., Chem. Commun., **1985**, 602-603 (1985)
430. N. Fujii, S. Futaki, K. Akaji, H. Yajima, A. Inoue & T. Segawa
Application of *N*^m-Mesitylenesulfonyl Tryptophan for the Synthesis of Neuromedin B and Neuromedin C.
Chem. Pharm. Bull., **33**, 3731-3737 (1985)
431. N. Fujii, H. Hayashi, S. Katakura, K. Akaji, H. Yajima, A. Inoue & T. Segawa
Application of New Heterobifunctional Crosslinking Reagents for the Preparation of Neurokinin (A and B)-BSA(Bovine Serum Albumin) Conjugates.
Int. J. Peptide Protein Res., **26**, 121-129 (1985)
432. R. Hosotani, K. Inoue, N. Fujii, H. Yajima & T. Tobe
Effect of Synthetic Neuromedin C, a Decapeptide of Gastrin Releasing Peptide ((GRP)[18-28]), on Blood Flow and Exocrine Secretion on the Pancreas in Dogs.
Life Science., **36**, 2429-2434 (1985)
433. H. Yajima, N. Fujii, K. Akaji, M. Sakurai, M. Nomizu, K. Mizuta, M. Aono, M. Moriga, K. Inoue, R. Hosotani & T. Tobe
Synthesis of a 42-Residue Peptide Corresponding to the Entire Amino Acid Sequence of Human GIP.
Chem. Pharm. Bull., **33**, 3578-3581 (1985)
434. K. Akaji, N. Fujii & H. Yajima
Synthesis of the Protected Tetracosapeptide Corresponding to Positions 30-53 of Mouse Epidermal Growth Factor (EGF).
Chem. Pharm. Bull., **33**, 173-183 (1985)
435. K. Akaji, N. Fujii, H. Yajima, K. Hayashi, k. Mizuta, M. Aono & M. Moriga
Synthesis of a Tripentacontapeptide with Epidermal Growth Factor Activity.

Chem. Pharm. Bull., **33**, 184-201 (1985)

436. S. Kiyama, K. Kitagawa, T. Akita, W.Y. Chey, A. Ayalp, K. Otsuki, S. Funakoshi, N. Fujii & H. Yajima
Alternative Synthesis of Porcine Secretin and Apparent Autolysis of the Product.
Chem. Pharm. Bull., **33**, 3205-3217 (1985)
437. H. Yajima, S. Futaki, N. Fujii, K. Akaji, S. Funakoshi, M. Sakurai, S. Katakura, K. Inoue, R. Hosotani, T. Tobe, T. Segawa, A. Inoue, K. Tatemoto & V. Mutt
Synthesis of Galanin, a New Gastrointestinal Polypeptide.
J. Chem. Soc. Chem. Commun., **1985**, 877-878 (1985)
438. N. Fujii, S. Futaki, K. Yasumura & H. Yajima
Mn-Mesitylenesulfonyltryptophan, a New Derivative for Peptide Synthesis.
Chem. Pharm. Bull., **32**, 2660-2665 (1984)
439. N. Fujii, M. Shimokura, K. Akaji, S. Kiyama & H. Yajima
Synthesis of the Protected Docosapeptide Corresponding to the C-Terminal Portion of Human Growth Hormone Releasing Factor (GRF, Somatocrinin).
Chem. Pharm. Bull., **32**, 510-519 (1984)
440. N. Fujii, M. Shimokura, M. Nomizu, H. Yajima, F. Shono, M. Tsuda & A. Yoshitake
Solution Synthesis of the Tetratetracontapeptide Amide Corresponding to the Entire Amino Acid Sequence of Growth Hormone Releasing Factor, Somatocrinin.
Chem. Pharm. Bull., **32**, 520-529 (1984)
441. N. Fujii, W. Lee, M. Shimokura & H. Yajima
Synthesis of a Hybrid Growth Hormone Releasing Factor (GRF)-PHI Heptacosapeptide Amide.
Chem. Pharm. Bull., **32**, 739-743 (1984)
442. N. Fujii, M. Shimokura, H. Yajima, F. Shono, M. Tsuda & A. Yoshitake
Synthesis of Growth Hormone Releasing Factor (GRF-37-NH₂) and N^a-Biotinyl GRF-44-NH₂.
Chem. Pharm. Bull., **32**, 1200-1208 (1984)
443. N. Fujii, M. Shimokura, H. Yajima, R. Shono, M. Tsuda & A. Yoshitake
Synthesis of Growth Hormone Releasing Factor (hGRF-40-OH).
Chem. Pharm. Bull., **32**, 1193-1199 (1984)
444. S. Kiyama, N. Fujii, H. Yajima, M. Moriga & A. Takagi
Alternative Synthesis of the Heptacosapeptide Amide Corresponding to the Entire Amino Acid Sequence of Chicken Secretin.

Int. J. Peptide Protein Res., **23**, 174-186 (1984)

445. H. Yajima, N. Fujii, M. Shimokura, K. Akaji, S. Kiyama & M. Nomizu
Solution Synthesis of a Tetratetracontapeptide Amide with Growth Hormone Releasing Activity.
Chem. Pharm. Bull., **31**, 1800-1802 (1983)
446. Takagi, M. Moriga, H. Uchino, K. Akaji, N. Fujii & H. Yajima
Distribution of Gastrin-Releasing Peptide in Mammalian Tissue.
Gut Peptide and Ulcer., **1983**, 33-37 (1983)
447. N. Nishi, A. Tsutsumi, M. Morishige, S. Kiyama, N. Fujii, M. Takeyama & H. Yajima
Apparent Autolysis of the *N*-Terminal Tetrapeptide of Vasoactive Intestinal Polypeptide (VIP).
Chem. Pharm. Bull., **31**, 1067-1072 (1983)
448. N. Fujii, W. Lee, H. Yajima, M. Moriga & K. Mizuta
Alternative Synthesis of Heptacosapeptide, a New Gastrointestinal Polypeptide.
Chem. Pharm. Bull., **31**, 3503-3514 (1983)
449. N. Fujii, S. Katakura, H. Yajima, Y. Nakata, A. Inoue & T. Segawa
Synthesis of Hylambates-Kassinin, a Frog Skin Tachykinin Peptide.
Chem. Pharm. Bull., **31**, 4259-4262 (1983)
450. K. Akaji, N. Fujii, H. Yajima, M. Moriga, A. Takagi, K. Mizuta, M. Noguchi & T.J. McDonald
Synthesis of Chicken Gastrin-Releasing Peptide.
Int. J. Peptide Protein Res., **20**, 276-288 (1982)
451. H. Yajima, Y. Minamitake, S. Funakoshi, I. Kitamura, N. Fujii, T. Segawa, Y. Nakata, T. Yasuhara & T. Nakajima
Synthesis of a Physalaemin-like Peptide, [Lys5, Thr6]-Physalaemin, Isolated from the Skin of a Frog, *Uperolela Rugosa*.
Chem. Pharm. Bull., **30**, 344-348 (1982)
452. S. Funakoshi, N. Fujii, H. Yajima, C. Shigeno, I. Yamamoto, R. Morita & K. Torizuka
Synthesis of the Octatriacontapeptide Corresponding to Positions 1 to 38 of Human Parathyroid Hormone.
Chem. Pharm. Bull., **30**, 1706-1717 (1982)
453. K. Akaji, N. Fujii, H. Yajima & D. Pearson
Synthesis of Urotensin II.
Chem. Pharm. Bull., **30**, 349-353 (1982)

454. K. Akaji, N. Fujii, H. Yajima, M. Moriga, M. Aono & A. Takagi
Synthesis of a Heptacosapeptide Amide Corresponding to the Entire Amino Acid Sequence of Gastrin-Releasing Peptide.
Int. J. Peptide Protein Res., **18**, 180-194 (1981)
455. N. Fujii & H. Yajima
Total Synthesis of Bovine Pancreatic Ribonuclease A . Part 1. Synthesis of the Protected Pentadecapeptide Ester (Positions 110-124).
J. Chem. Soc. Perkin I., **1981**, 789-796 (1981)
456. N. Fujii & H. Yajima
Total Synthesis of Bovine Pancreatic Ribonuclease A. Par 2. Synthesis of the Protected Hexatriacontapeptide Ester (Positions 89-124).
J. Chem. Soc. Perkin I., **1981**, 797-803 (1981)
457. N. Fujii & H. Yajima
Total Synthesisof Bovine Pancreatic Ribonuclease A. Part 3. Synthesis of the Protected Hexapentacontapeptide Ester (Positions 69-124).
J. Chem. Soc. Perkin I., **1981**, 804-810 (1981)
458. N. Fujii & H. Yajima
Total Synthesis of Bovine Pancreatic Ribonuclease A. Part 4. Synthesis of the Protected Tetraoctacontapeptide Ester (Positions 41-124).
J. Chem. Soc. Perkin I., **1981**, 811-818 (1981)
459. N. Fujii & H. Yajima
Total Syhnthesis of Bovine Pancreatic Ribonuclease A. Part 5. Synthesis of the Protected S-Protein (Positions 21-124) and the Protected S-Peptide (Positions 1-20).
J. Chem. Soc. Perkin I., **1981**, 819-830 (1981)
460. N. Fujii & H. Yajima
Total Synthesis of Bovine Pancreatic Ribonuclease A. Part 6. Synthesis of RNase A with Full Enzymic Activity.
J. Chem. Soc. Perkin I., **1981**, 831-841 (1981)
461. H. Yajima & N. Fujii
Chemical Synthesis of Crystalline Ribonuclease A.
Chem. Pharm. Bull., **29**, 600-602 (1981)
462. H. Yajima & N. Fujii
Chemical Synthesisof a Crystalline Protein with the Full Enzymatic Activity of Ribonuclease A.
J. Am. Chem. Soc., **103**, 5867-5871 (1981)

463. H. Yajima & N. Fujii
Totally Synthetic Crystalline Ribonuclease A.
Biopolymers, **20**, 1859-1867 (1981)
464. N. Fujii & H. Yajima
Synthesis of Ribonuclease S'.
Chem. Pharm. Bull., **29**, 1927-1934 (1981)
465. H. Irie, A. Shiina, T. Fushimi, J. Katakawa, N. Fujii & H. Yajima
New Synthesis of Isoquinoline Alkaloids, Thalifoline, Corypalline and Cherylline.
Chem. Lett., **1980**, 875-878 (1980)
466. H. Irie, H. Nakanishi, N. Fujii, Y. Mizuno, T. Fushimi, S. Funakoshi & H. Yajima
Validity of Methoxycarbonyl as an *N*-Protecting Group on Peptide Synthesis.
New Synthesis of MSH-Release Inhibiting Factor.
Chem. Lett., **1980**, 705-708 (1980)
467. H. Yajima, K. Akaji, Y. Hirota & N. Fujii
Thiazolidine-2-thione as a Carboxyl-Activating Reagent for Peptide Synthesis.
Chem. Pharm. Bull., **28**, 3140-3142 (1980)
468. H. Yajima, K. Akaji, S. Funakoshi, N. Fujii & H. Irie
Behaviour of *S*-Acetamidomethylcysteine Sulfoxide under Deprotecting Conditions in Peptide Synthesis.
Chem. Pharm. Bull., **28**, 1942-1945 (1980)
469. S. Funakoshi, N. Fujii & H. Yajima
A New Deprotecting Procedure for *p*-Toluenesulfonyl and *p*-Methoxybenzenesulfonyl Groups from the *N*^m-Function of Histidine.
Chem. Pharm. Bull., **28**, 962-931 (1980)
470. H. Yajima, K. Akaji, N. Fujii, M. Moriga, M. Aono & A. Takagi
Synthesis of a Heptacosapeptide Amide Corresponding to the Entire Amino Acid Sequence of Gastrin Releasing Peptide.
Chem. Pharm. Bull., **28**, 2276-2278 (1980).
471. H. Yajima & N. Fujii
Chemical Synthesis of Bovine Pancreatic Ribonuclease A.
J. Chem. Soc., Chem. Commun., **1980**, 115-116 (1980)
472. H. Yajima, K. Akaji, K. Mitani, N. Fujii, S. Funakoshi, H. Adachi, M. Oishi & Y. Akazawa
Application of a New Arginine Derivative, *N*^G-Mesitylene-2-sulfonylarginine, to

- the Synthesis of Substance P and Neurotensin.
Int. J. Peptide Protein Res., **14**, 169-176 (1979)
473. K. Kitagawa, K. Kitada, Y. Kiso, T. Akita, S. Funakoshi, N. Fujii & H. Yajima
N^m-p-Methoxyphenylsulfonylhistidine, a New Derivative for Peptide Synthesis.
J. Chem.Soc. Chem. Commun., **1979**, 955-956 (1979)
474. H. Yajima, S. Funakoshi, N. Fujii, K. Akaji & H. Irie
Behaviour of *S*-Substituted Cysteine Sulfoxide under Acidolytical Deprotecting Conditions.
Chem. Pharm. Bull., **27**, 1060-1061 (1979)
475. S. Funakoshi, N. Fujii, k. Akaji, H. Irie & H. Yajima
Behabiour of *S*-Substituted Cysteine Sulfoxide under Deprotecting Conditions in Peptide Synthesis.
Chem. Pharm. Bull., **27**, 2151-2156 (1979)
476. H. Yajima, T. Sasaki, H. Ogawa, N. Fujii, T. Segawa & Y. Nakata
Synthesis of Kassininm, a New Frog Skin Peptide.
Chem. Pharm. Bull., **26**, 1231-1235 (1978)
477. N. Fujii, T. Sasaki, S. Funakoshi, H. Irie & H. Yajima
Convenient Procedure for the Preparation of Methionine Sulfoxide Derivatives.
Chem. Pharm. Bull., **26**, 650-653 (1978).
478. H. Irie, N. Fujii, H. Ogawa, H. Yajima, M. Fujino & S. Shinagawa Examination of the Methanesulfonic Acid Procedure for the Synthesis of Peptides Containing Methioine.
Chem. Pharm. Bull., **25**, 2929-2934 (1977)
479. N. Fujii, H. Irie & H. Yajima
Regioselective Cleavage of Aromatic Methyl Ether by Methanesulfonic Acid in the Presence of Methionine.
J. Chem. Soc. Perkin I., **1977**, 2288-2289 (1977)
480. H. Yajima, H. Ogawa, N. Fujii & S. Funakoshi
Selective Removal of Acid Labile α -Amino Protecting Groups with Dilute Sulfonic Acids.
Chem. Pharm. Bull., **25**, 740-747 (1977)
481. N. Fujii, S. Funakoshi, T. Sasaki & H. Yajima
Examination of the *N*-Alkylation of Lysine in the Methanesulfonic Acid Procedure for Peptide Synthesis.
Chem. Pharm. Bull., **25**, 3096-3098 (1977)
482. F. Tamura, H. Ogawa, N. Fujii, H. Yajima, K. Miyata, N. Nakamura & A.

Tanaka

Synthesis of Two Octadecapeptides Corresponding to the Entire Amino Acid Sequence of Camel β -MSH.

F. Tamura, H. Ogawa, N. Fujii, H. Yajima, K. Miyata, N. Nakamura & A. Tanaka
Chem. Pharm. Bull., **25**, 767-774 (1977)

483. K. Kitagawa, T. Akita, T. Segawa, M. Nakano, N. Fujii & H. Yajima
Conventional Synthesis of a New Hypothalamic Peptide, Bovine Neurotensin.
Chem. Pharm. Bull., **24**, 2692-2698 (1976)

484. H. Irie, N. Fujii, H. Ogawa & H. Yajima
Role of Met in the Facilitated Cleavage of Aromatic Ethers by Methanesulphonic Acid.

J. Chem. Soc., Chem. Commun., **1976**, 922-923 (1976)

485. H. Yajima, M. Kurobe, I. Yo, N. Fujii & Y. Baba
Applicaion of the Solid Phase Synthesis of the Preparation of Pro-Analogues of LH and FSH Releasing Hormone.
Chem. Pharm. Bull., **23**, 1622-1624 (1975)

486. H. Yajima, H. Ogawa, H. Watanabe, N. Fujii, M. Kurobe & S. Miyamoto
Application of the Trifluoromethanesulfonic Acid Procedure to the Synthesis of Tuftsin.

Chem. Pharm. Bull., **23**, 371-371 (1975)

487. H. Yajima, Y. Kiso, h. Ogawa, N. Fujii & H. Irie
Acidolysis of Protecting Groups in Peptide Chemistry by Fluorosulphonic Acid and Methanesulphonic Acid.

Chem. Pharm. Bull., **23**, 1164-1166 (1975)

488. N. Fujii & H. Yajima
Application of the Methanesulfonic Acid Procedure to the Synthesis of a Peptide with Somatostatin Activity.

Chem. Pharm. Bull., **23**, 1596-1603 (1975)

489. N. Fujii & H. Yajima
Synthesis of N¹-Tyr-Somatostatin.
Chem. Pharm. Bull., **23**, 2446-2499 (1975)

490. H. Yajima, N. Fujii, H. Ogawa & H. Kawatani
Trifluoromethanesulfonic Acid, as a Deprotecting Reagent in Peptide Chemistry
J. Chem. Soc., Chem. Commun., **1974**, 107-108 (1974).

総説

1. 大石真也、藤井信孝
ケモカイン受容体 CXCR4 とそのアンタゴニスト.
和光純薬時報. **78**(1) 2-5. (2010)
2. Hyunsuk S., Oishi S., Fujii N.
Chemokine receptor CXCR4 as a therapeutic target for neuroectodermal tumors.
Semin. Cancer Biol. **9**(2) 123-134 (2009)
Tamamura H., Tsutsumi H., Nomura W., Tanaka T., Fujii N.
A future perspective on the development of chemokine receptor CXCR4 antagonists.
Expert Opin. Drug Discov. **3**(10) 1155-1166 (2008)
3. Oishi S., Narumi T., Ohno H., Otaka A., Fujii N.
Synthesis of highly functionalized alkene dipeptide isosteres and its application to the structure-activity relationship study on bioactive peptides.
J. Synth. Org. Chem. Jpn. **66**(9) in press (2008)
4. Tamamura H., Tsutsumi H., Masuno H., Fujii N.
Development of low molecular weight CXCR4 antagonists by exploratory structural tuning of cyclic tetra- and pentapeptide-scaffolds towards the treatment of HIV infection, cancer metastasis and rheumatoid arthritis.
Curr. Med. Chem. **14**(1) 93-102 (2007)
5. Tsutsumi H., Tanaka T., Ohashi N., Masuno H., Tamamura H., Hiramatsu K., Araki T., Ueda S., Oishi S., Fujii N.
The therapeutic potential of the chemokine receptor CXCR4 antagonists as multi-functional agents.
Biopolymers, **88**(2) 279-289 (2007)
6. Tsutsumi H., Tamamura H., Fujii N.
Inhibitors of the chemokine receptor CXCR4: chemotherapy of AIDS, metastatic cancer, leukemia and rheumatoid arthritis.
Lett. Drug Design Discov. **4**(1) 20-26 (2007)
7. 大石真也、藤井信孝
ペプチドをリードとした創薬
Tanpakushitsu Kakusan Koso, **52**(13,zokan) 1696-1701 (2007)
8. 大石真也、藤井信孝

ケミカルバイオロジーを基盤とするペプチド創薬

Idenshi Igaku Mook, **8** 87-91 (2007)

9. Tamamura H., Tsutumi H., Fujii N.
The chemokine receptor CXCR4 as a therapeutic target for several diseases.
Mini Rev. Med. Chem. **6**(9) 989-995 (2006)
10. Otaka A., Fujii N.
Development of antiviral fusion inhibiting peptides.
Kagaku Kogyo, **57**(10) 797-801 (2006)
11. Tamamura H., Fujii N.
The therapeutic potential of CXCR4 antagonists in the treatment of HIV infection, cancer metastasis and rheumatoid arthritis.
Expert Opin. Ther. Targets, **9**(6) 1267-1282 (2005)
12. Fujii N., Otaka A., Tamamura H.
Innovative platform for drug discovery based on chemical proteomics: from knowledge to controlling.
Saibo Kogaku **24**(11) 1181-1186 (2005)
13. Tamamura H., Otaka A., Fujii N.
Development of anti-HIV agents targeting dynamic supramolecular mechanism: entry and fusion inhibitors based on CXCR4/CCR5 antagonists and gp41-C34-remodeling peptides.
Curr. HIV Res. **3**(4) 289-301 (2005)
14. Tamamura H., Fujii N.
Two orthogonal approaches to overcome multi-drug resistant HIV-1s: development of protease inhibitors and entry inhibitors based on CXCR4 antagonists.
Curr. Drug Targets Infec. Disord. **4**(2) 103-110 (2004)
15. Otaka A., Mitsuyama E., Watanabe J., Watanabe H., Fujii N.
Synthesis of fluorine-containing bioisosteres corresponding to phosphoamino acids and dipeptide units.
Biopolymers, **76**(2) 140-149 (2004)
16. Fujii N.
Medicinal chemistry based on proteomics: from knowledge to controlling.
BIO. Clinica. **19**(14) 1178-1184 (2004)
17. Otaka A., Fujii N., Yamamoto N.
Prospect of SARS drugs.
Gendai Iryo, **36**(11) 2198-2203 (2004)

18. Fujii N., Nakashima H., Tamamura H.
The therapeutic potential of CXCR4 antagonists in the treatment of HIV.
Expert Opinion on Investigational Drugs, **12**(2) 185-195 (2003)
19. Fujii N.
Genomic information-based astringent drug discovery and development.
バイオベンチャー (5) 24-29 (2002)

著書

1. インシリコ創薬科学-ゲノム情報から創薬へ- (共著 京都廣川書店 2008)
2. 新生化学実験講座 第1巻. タンパク質 VI. 合成および発現 (共著 東京化学同人、1992)
3. 新生化学実験講座 第9巻. ホルモン. ペプチド合成: 下垂体ホルモンの合成 (共著 東京化学同人 1991)
4. 医薬品の開発第3巻. 生体成分、ホルモン系ペプチド、タンパク質: 視床下部、下垂体以外のペクチドタンパク質ホルモン(共著 廣川書店、1989)