

## 原著論文（主要なもの）

- 1 Yoshimitsu, Y.; Oishi, S.; Miyagaki, J.; Inuki, S.; Ohno, H.; Fujii, N. Pachastrissamine (jaspine B) and its stereoisomers inhibit sphingosine kinases and atypical protein kinase C. *Bioorg. Med. Chem.* **2011**, 19(18) 5402-5408.
- 2 Takeuchi, T.; Oishi, S.; Watanabe, T.; Ohno, H.; Sawada, J.; Matsuno, K.; Asai, A.; Asada, N.; Kitaura, K.; Fujii, N. Structure-activity relationships of carboline and carbazole derivatives as a novel class of ATP-competitive kinesin spindle protein inhibitors. *J. Med. Chem.* **2011**, 54(13) 4839-4846.
- 3 Inokuchi, E.; Oishi, S.; Kubo, T.; Ohno, H.; Shimura, K.; Matsuoka, M.; Fujii, N. Potent CXCR4 antagonists containing amidine-type peptide bond isosteres. *ACS Med. Chem. Lett.* **2011**, 2(6) 477-480.
- 4 Masuda, R.; Oishi, S.; Ohno, H.; Kimura, H.; Saji, H.; Fujii, N. Concise site-specific synthesis of DTPA-peptide conjugates: application to imaging probes for the chemokine receptor CXCR4. *Bioorg. Med. Chem.* **2011**, 19(10) 3216-3220.
- 5 Oishi, S.; Misu, R.; Tomita, K.; Setsuda, S.; Masuda, R.; Ohno, H.; Naniwa, Y.; Ieda, N.; Inoue, N.; Ohkura, S.; Uenoyama, Y.; Tsukamura, H.; Maeda, K.; Hirasawa, A.; Tsujimoto, G.; Fujii, N. Activation of neuropeptide FF receptors by kisspeptin receptor ligands. *ACS Med. Chem. Lett.* **2011**, 2(1) 53-57.
- 6 Oishi, S.; Watanabe, K.; Ito, S.; Tanaka, M.; Nishikawa, H.; Ohno, H.; Shimane, K.; Izumi, K.; Sakagami, Y.; Kodama, E. N.; Matsuoka, M.; Asai, A.; Fujii, N. Affinity selection and sequence-activity relationships of HIV-1 membrane fusion inhibitors directed at the drug-resistant variants. *Med. Chem. Commun.* **2010**, 1(4) 276-281.
- 7 Oishi, S.; Watanabe, T.; Sawada, J.; Asai, A.; Ohno, H.; Fujii, N. Kinesin spindle protein (KSP) inhibitors with 2,3-fused indole scaffolds. *J. Med. Chem.* **2010**, 53(13) 5054-5058.
- 8 Narumi, T.; Hayashi, R.; Tomita, K.; Kobayashi, K.; Tanahara, N.; Ohno, H.; Naito, T.; Kodama, E.; Matsuoka, M.; Oishi, S.; Fujii, N. Synthesis and biological evaluation of selective CXCR4 antagonists containing alkene dipeptide isosteres. *Org. Biomol. Chem.* **2010**, 8(3) 616-621.
- 9 Kajiwara, K.; Watanabe, K.; Tokiwa, R.; Kurose, T.; Ohno, H.; Tsutsumi, H.; Hata, Y.; Izumi, K.; Kodama, E.; Matsuoka, M.; Oishi, S.; Fujii, N. Bioorganic synthesis of a recombinant HIV-1 fusion inhibitor, SC35EK, with an N-terminal pyroglutamate capping group. *Bioorg. Med. Chem.* **2009**, 17(23) 7964-7970.
- 10 Tanaka, M.; Kajiwara, K.; Tokiwa, R.; Watanabe, K.; Ohno, H.; Tsutsumi, H.; Hata, Y.; Izumi, K.; Kodama, E.; Matsuoka, M.; Oishi, S.; Fujii, N. Bioorganic synthesis of end-capped anti-HIV peptides by simultaneous cyanocysteine-mediated cleavages of recombinant proteins. *Bioorg. Med. Chem.* **2009**, 17(21) 7487-7492.
- 11 Watabe, T.; Terakawa, Y.; Watanabe, K.; Ohno, H.; Nakano, H.; Nakatsu, T.; Kato, H.; Izumi, K.; Kodama, E.; Matsuoka, M.; Kitaura, K.; Oishi, S.; Fujii, N. X-ray crystallographic study of an HIV-1 fusion inhibitor with the gp41 S138A substitution. *J. Mol. Biol.* **2009**, 392(3) 657-665.
- 12 Oishi, S.; Kodera, Y.; Nishikawa, H.; Kamitani, H.; Watabe, T.; Ohno, H.; Tochikura, T.; Shimane, K.; Kodama, E.; Matsuoka, M.; Mizukoshi, F.; Tsujimoto, H.; Fujii, N. Design and synthesis of membrane fusion inhibitors against the feline immunodeficiency virus. *Bioorg. Med. Chem.* **2009**, 17(14) 4916-4920.
- 13 Oishi, S.; Kamitani, H.; Kodera, Y.; Watanabe, K.; Kobayashi, K.; Narumi, T.; Tomita, K.; Ohno, H.; Naito, T.; Kodama, E.; Matsuoka, M.; Fujii, N. Peptide bond mimicry by (*E*)-alkene and (*Z*)-fluoroalkene peptide isosteres: synthesis and bioevaluation of  $\alpha$ -helical anti-HIV peptide analogues. *Org. Biomol. Chem.* **2009**, 7(14) 2872-2877.
- 14 Tomita, K.; Popiel, H.A.; Nagai, Y.; Toda, T.; Yoshimitsu, Y.; Ohno, H.; Oishi, S.; Fujii, N. Structure-activity relationship study on polyglutamine binding peptide QBP1. *Bioorg. Med. Chem.* **2009**, 17(3) 1259-1263.
- 15 Tomita, K.; Oishi, S.; Ohno, H.; Peiper, S.C.; Fujii, N. Development of novel G protein-coupled receptor 54 agonists with resistance to degradation by matrix metalloproteinase. *J. Med. Chem.* **2008**, 51(23) 7645-7649.
- 16 Nishikawa H.; Oishi S.; Fujita M.; Watanabe K.; Tokiwa R.; Ohno H.; Kodama E.; Izumi K.; Kajiwara K.; Naitoh T.; Matsuoka M.; Otaka A.; Fujii N. Identification of minimal sequence for HIV-1 fusion inhibitors. *Bioorg. Med. Chem.* **2008**, 16(20), 9184-9187.

- 17 Ueda, S.; Kato, M.; Ohno, H.; Evans, B.; Wang, Z.; Peiper, S. C.; Izumi, K.; Kodama, E.; Matsuoka, M.; Nagasawa, H.; Oishi, S.; Fujii, N. Identification of novel nonpeptide CXCR4 antagonists by ligand-based design approach. *Bioorg. Med. Chem. Lett.* **2008**, 18(14) 4124-4129.
- 18 Tomita, K.; Oishi, S.; Ohno, H.; Fujii, N. Structure-activity relationship study and NMR analysis of fluorobenzoyl pentapeptide GPR54 agonists. *Biopolymers*, **2008**, 90(4) 503-511.
- 19 Oishi, S.; Masuda, R.; Evans, B.; Ueda, S.; Goto, Y.; Ohno, H.; Hirasawa, A.; Tsujimoto, G.; Wang, Z.; Peiper, S. C.; Naito, T.; Kodama, E.; Matsuoka, M.; Fujii, N. Synthesis and application of fluorescein- and biotin-labeled molecular probes for chemokine receptor CXCR4. *ChemBioChem*, **2008**, 9(7) 1154-1158.
- 20 Oishi, S.; Ito, S.; Nishikawa, H.; Watanabe, K.; Tanaka, M.; Ohno, H.; Izumi, K.; Sakagami, Y.; Kodama, E.; Matsuoka, M.; Fujii, N. Design of a novel HIV-1 fusion inhibitor that displays a minimal interface for binding affinity. *J. Med. Chem.* **2008**, 51(3) 388-391.
- 21 Tomita, K.; Oishi, S.; Cluzeau, J.; Ohno, H.; Navenot, J.-M.; Wang, A.; Peiper, S. C.; Akamatsu, M.; Fujii, N. SAR and QSAR studies on the N-terminally acylated pentapeptide agonists for GPR54. *J. Med. Chem.* **2007**, 50(14) 3222-3228.
- 22 Cluzeau, J.; Oishi, S.; Ohno, H.; Wang, Z.; Evans, B.; Peiper, S. C.; Fujii, N. Design and synthesis of all diastereomers of cyclic pseudo-dipeptides as mimic of cyclic CXCR4 pentapeptide antagonist. *Org. Biomol. Chem.* **2007**, 5(12) 1915-1923.
- 23 Tomita, K.; Narumi, T.; Niida, A.; Oishi, S.; Ohno, H.; Fujii, N. Fmoc-based solid-phase synthesis of GPR54-agonistic pentapeptide derivatives containing alkene- and fluoroalkene-dipeptide isosteres. *Biopolymers*, **2007**, 88(2) 272-278.
- 24 Tanaka, M.; Oishi, S.; Ohno, H.; Fujii, N. A novel oxazolidine linker for the synthesis of peptide aldehydes. *Int. J. Pep. Res. Ther.* **2007**, 13(1-2) 271-279.
- 25 Ueda, S.; Oishi, S.; Wang, Z.-x.; Araki, T.; Tamamura, H.; Cluzeau, J.; Ohno, H.; Kusano, S.; Nakashima, H.; Trent, J.O.; Peiper, S.C.; Fujii, N. Structure-activity relationships of cyclic peptide-based chemokine receptor CXCR4 antagonists: disclosing the importance of side-chain and backbone functionalities. *J. Med. Chem.* **2007**, 50(2) 192-198.
- 26 Tomita, K.; Niida, A.; Oishi, S.; Ohno, H.; Cluzeau, J.; Navenot, J.-M.; Wang, Z.; Peiper, S.C.; Fujii, N. Structure-activity relationship study on small peptidic GPR54 agonists. *Bioorg. Med. Chem.* **2006**, 14(22) 7595-7603.
- 27 Oishi, S.; Miyamoto, K.; Niida, A.; Yamamoto, M.; Ajito, K.; Tamamura, H.; Otaka, A.; Kuroda, Y.; Asai, A.; Fujii, N. Application of tri- and tetrasubstituted alkene dipeptide mimetics to conformational studies of cyclic RGD peptides. *Tetrahedron*, **2006**, 62(7) 1416-1424.
- 28 Oishi, S; Karki, R. G.; Shi, Z-D.; Worthy, K. M.; Bindu, L.; Chertov, O.; Esposito, D.; Frank, P.; Gillette, W. K.; Maderia, M. A.; Hartley, J.; Nicklaus, M. C.; Barchi, J. J., Jr.; Fisher, R. J.; Burke, T. R., Jr. Evaluation of macrocyclic Grb2 SH2 domain-binding peptide mimetics prepared by ring-closing metathesis of C-terminal allylglycines with an N-terminal  $\beta$ -vinyl-substituted phosphotyrosyl mimetic. *Bioorg. Med. Chem.* **2005**, 13(7) 2431-2438.
- 29 Oishi, S.; Karki, R. G.; Kang, S-U.; Wang, X.; Worthy, K. M.; Bindu, L. K.; Nicklaus, M. C.; Fisher, R. J.; Burke, T. R., Jr. Design and synthesis of conformationally constrained Grb2 SH2 domain binding peptides employing  $\gamma$ -methylphenylalanyl based phosphotyrosyl mimetics. *J. Med. Chem.* **2005**, 48(3) 764-772.
- 30 Oishi, S.; Shi, Z.-D.; Worthy, K. M.; Bindu, L. K.; Fisher, R. J.; Burke, T. R. Jr. Ring-closing metathesis of C-terminal allylglycine residues with an N-terminal  $\beta$ -vinyl-substituted phosphotyrosyl mimetic as an approach to novel Grb2 SH2 domain-binding macrocycles. *ChemBioChem*, **2005**, 6(4) 668-674.
- 31 Oishi, S.; Kang, S.-K.; Liu, H.; Zhang, M.; Yang, D.; Deschamps, J. R.; Burke, T. R., Jr. Synthesis of  $\alpha,\alpha$ -disubstituted 4-phosphonophenylalanine analogues as conformationally-constrained phosphotyrosyl mimetics. *Tetrahedron*, **2004**, 60(13) 2971-2977.
- 32 Fujii, N.; Oishi, S.; Hiramatsu, K.; Araki, T.; Ueda, S.; Tamamura, H.; Otaka, A.; Kusano, S.; Terakubo, S.; Nakashima, H.; Broach, J. A.; Trent, J. O.; Wang, Z.; Peiper, S. C. Molecular-size reduction of a potent CXCR4-chemokine antagonist using orthogonal combination of conformation- and sequence-based libraries. *Angew. Chem. Int. Ed. Engl.* **2003**, 42(28) 3251-3253.

- 33 Oishi, S.; Kamano, T.; Niida, A.; Odagaki, Y.; Hamanaka, H.; Yamamoto, M.; Ajito, K.; Tamamura, H.; Otaka, A.; Fujii, N. Diastereoselective synthesis of new  $\gamma$ [(E)-CH=CMe]- and  $\gamma$ [(Z)-CH=CMe]-type alkene dipeptide isosteres by organocopper reagents and application to conformationally restricted cyclic RGD peptidomimetics. *J. Org. Chem.* **2002**, 67(17) 6162-6173.
- 34 Oishi, S.; Niida, A.; Kamano, T.; Odagaki, Y.; Hamanaka, N.; Yamamoto, M.; Ajito, K.; Tamamura, H.; Otaka, A.; Fujii, N. Regio- and stereoselective ring-opening of chiral 1,3-oxazolidin-2-one derivatives by organocopper reagents provides novel access to di-, tri- and tetra-substituted alkene dipeptide isosteres. *J. Chem. Soc., Perkin Trans. 1*, **2002**, (15) 1786-1793.
- 35 Oishi, S.; Kamano, T.; Niida, A.; Odagaki, Y.; Tamamura, H.; Otaka, A.; Hamanaka, H.; Fujii, N. Diastereoselective synthesis of  $\gamma$ [(E)-CH=CMe]- and  $\gamma$ [(Z)-CH=CMe]-type dipeptide isosteres by organocopper-mediated anti-S<sub>N</sub>2' reaction. *Org. Lett.* **2002**, 4(7) 1051-1054.
- 36 Oishi, S.; Niida, A.; Kamano, T.; Odagaki, Y.; Tamamura, H.; Otaka, A.; Hamanaka, H.; Fujii, N. Diastereoselective synthesis of  $\gamma$ [(E)-CMe=CH]- and  $\gamma$ [(E)-CMe=CMe]-type dipeptide isosteres based on organocopper-mediated anti-S<sub>N</sub>2' reaction. *Org. Lett.* **2002**, 4(7) 1055-1058.
- 37 Oishi, S.; Tamamura, H.; Yamashita, M.; Odagaki, Y.; Hamanaka, N.; Otaka, A.; Fujii, N. Stereoselective synthesis of a set of two functionalized (E)-alkene dipeptide isosteres of L-amino acid-L-Glu and L-amino acid-D-Glu. *J. Chem. Soc., Perkin Trans. 1*, **2001**, (19) 2445-2451.

### 原著論文（その他）

- 38 Inoue, N.; Tomikawa, J.; Sasaki, Y.; Uchida, A.; Oishi, S.; Fujii, N.; Uenoyama, Y.; Yamamoto, N.; Maeda, K.; Tsukamura, H. Identification of suncus kisspeptin and induction of ovulation by peripheral kisspeptin administration in the musk shrew (*Suncus murinus*) reflex ovulator. *Proc. Natl. Acad. Sci.* **2011**, 108(42) 17527-17532.
- 39 Hirano, K.; Inaba, Y.; Takasu, K.; Oishi, S.; Takemoto, Y.; Fujii, N.; Ohno H. Gold(I)-catalyzed polycyclizations of polyenyne-type anilines based on hydroamination and consecutive hydroarylation cascade. *J. Org. Chem.* **2011**, 76(21) 9068-9080.
- 40 Nishizawa, K.; Nishiyama, H.; Matsui, Y.; Kobayashi, T.; Kotani, H.; Masutani, H.; Oishi, S.; Saito, R.; Toda, Y.; Fujii, N.; Yodoi, J.; Ogawa, O. Thioredoxin interacting protein suppresses bladder carcinogenesis. *Carcinogenesis*, **2011**, 32(10) 1459-1466.
- 41 Izumi, K.; Watanabe, K.; Oishi, S.; Fujii, N.; Matsuoka, M.; Sarafianosc, S.G.; Kodama, E. Potent anti-HIV-1 activity of N-HR-derived peptides including a deep pocket-forming region without antagonistic effect on T-20. *Antivir. Chem. Chemother.* **2011**, 22(1) 51-55
- 42 Iwata, A.; Inuki, S.; Oishi, S.; Fujii, N.; Ohno, H. Formal total synthesis of (+)-lysergic acid via zinc(II)-mediated regioselective ring-opening reduction of 2-alkynyl-3-indolyloxirane. *J. Org. Chem.* **2011**, 76(13) 5506-5512.
- 43 Tokimizu, Y.; Ohta, Y.; Chiba, H.; Oishi, S.; Fujii, N.; Ohno, H. Direct synthesis of highly fused perimidines by copper(I)-catalyzed hydroamination of 2-ethynylbenzaldehydes. *Tetrahedron*, **2011**, 67(29) 5168-5175.
- 44 Kuil, J.; Yuan, H.; Buckle, T.; van den Berg, N. S.; Oishi, S.; Fujii, N.; Josephson, L.; van Leeuwen, F. W. B. Synthesis and evaluation of a bimodal CXCR4 antagonistic peptide. *Bioconjug. Chem.* **2011**, 22(5) 859-864.
- 45 Dar, A.; Schajnovitz, A.; Lapid, K.; Kalinkovich, A.; Itkin, T.; Ludin, A.; Kao, W.; Battista, M.; Tesio, M.; Kollet, O.; Cohen, N. N.; Margalit, R.; Buss, E. C.; Baleux, F.; Oishi, S.; Fujii, N.; Larochelle, A.; Dunbar, C. E.; Broxmeyer, H. E.; Frenette, P. S.; Lapidot, T. Rapid mobilization of hematopoietic progenitors by AMD3100 and catecholamines is mediated by CXCR4-dependent SDF-1 release from bone marrow stromal cells. *Leukemia*, **2011**, 25(8) 1286-1296.
- 46 Inokuchi, E.; Yamada, A.; Hozumi, K.; Tomita, K.; Oishi, S.; Ohno, H.; Nomizu, M.; Fujii, N. Design and synthesis of amidine-type peptide bond isostere: application of nitrile oxide derivatives as active ester equivalents to peptide and peptidomimetics synthesis. *Org. Biomol. Chem.* **2011**, 9(9) 3421-3427.
- 47 Xing, N. D.; Ding, S. T.; Saito, R.; Nishizawa, K.; Kobayashi, T.; Inoue, T.; Oishi, S.; Fujii, N.; Lv, J. J.; Ogawa, O.; Nishiyama, H. A potent chemotherapeutic strategy in prostate cancer: S-(methoxytrityl)-l-cysteine, a novel Eg5 inhibitor. *Asian J. Androl.* **2011**, 13(2) 236-241.

- 48 Kang, H.S.; Baba, T.; Mandai, M.; Matsumura, N.; Hamanishi, J.; Kharma, B.; Kondoh, E.; Yoshioka, Y.; Oishi, S.; Fujii, N.; Murphy S. K.; Konishi, I. GPR54 is a target for suppression of metastasis in endometrial cancer. *Mol. Cancer Ther.* **2011**, 10(4) 580-590.
- 49 Inuki, S.; Iwata, A.; Oishi, S.; Fujii, N.; Ohno, H. Enantioselective total synthesis of (+)-lysergic acid, (+)-lysergol, and (+)-isolysergol by palladium-catalyzed domino cyclization of allenes bearing amino and bromoindolyl groups. *J. Org. Chem.* **2011**, 76(7) 2072-2083.
- 50 Hirano, K.; Inaba, Y.; Takahashi, N.; Shimano, M.; Oishi, S.; Fujii, N.; Ohno, H. Direct synthesis of fused-indoles by gold-catalyzed cascade cyclization of diynes. *J. Org. Chem.* **2011**, 76(5) 1212-1227.
- 51 Noda, M.; Omatsu, Y.; Sugiyama, T.; Oishi, S.; Fujii, N.; Nagasawa, T. CXCL12-CXCR4 chemokine signaling is essential for NK cell development in adult mice. *Blood*, **2011**, 117(2) 451-458.
- 52 Shimura, K.; Nameki, D.; Kajiwara, K.; Watanabe, K.; Sakagami, Y.; Oishi, S.; Fujii, N.; Matsuoka, M.; Sarafianos, S. G.; Kodama, E. Resistance profiles of novel electrostatically constrained HIV-1 fusion inhibitors. *J. Biol. Chem.* **2010**, 285(50) 39471-39480.
- 53 Gravel, S.; Malouf, C.; Boulais, P.E.; Berchiche, Y.A.; Oishi, S.; Fujii, N.; Leduc, R.; Sinnott, D.; Heveker, N. The peptidomimetic CXCR4 antagonist TC14012 recruits  $\beta$ -arrestin to CXCR7 roles of receptor domains. *J. Biol. Chem.* **2010**, 285(49) 37939-37943.
- 54 Ohta, Y.; Tokimizu, Y.; Oishi, S.; Fujii, N.; Ohno, H. Direct synthesis of quinazolines through copper-catalyzed reaction of aniline-derived benzamidines. *Org. Lett.* **2010**, 12(17) 3963-3965.
- 55 Ding, S.; Nishizawa, K.; Kobayashi, T.; Oishi, S.; Lv, J.; Fujii, N.; Ogawa, O.; Nishiyama, H. A Potent Chemotherapeutic strategy for bladder cancer: (S)-methoxytrityl-L-cysteine, a novel Eg5 inhibitor. *J. Urol.* **2010**, 183(7) 314-318.
- 56 Inuki, S.; Yoshimitsu, Y.; Oishi, S.; Fujii, N.; Ohno, H. Ring-construction/stereoselective functionalization cascade: total synthesis of pachastrissamine (jaspine B) through palladium-catalyzed bis-cyclization of propargyl chlorides and carbonates. *J. Org. Chem.* **2010**, 75(11) 3831-3842.
- 57 Izumi, K.; Nakamura, S.; Nakano, H.; Shimura, K.; Sakagami, Y.; Oishi, S.; Uchiyama, S.; Ohkubo, T.; Kobayashi, Y.; Fujii, N.; Matsuoka, M.; Kodama, E.N. Characterization of HIV-1 resistance to a fusion inhibitor, N36, derived from the gp41 amino terminal heptad repeat. *Antiviral Res.* **2010**, 87(2) 179-186.
- 58 Yoshimitsu, Y.; Inuki, S.; Oishi, S.; Fujii, N.; Ohno, H. Stereoselective divergent synthesis of four diastereomers of pachastrissamine (jaspine B). *J. Org. Chem.* **2010**, 75(11) 3843-3846.
- 59 Okano, A.; Oishi, S.; Tanaka, T.; Fujii, N.; Ohno, H. Construction of linked nitrogen heterocycles by palladium(0)-catalyzed intramolecular domino cyclization of 2-alkynylaziridines bearing a 2-aminoethyl group via ring expansion with isocyanate. *J. Org. Chem.* **2010**, 75(10) 3396-3400.
- 60 Okano, A.; Tsukamoto, K.; Kosaka, S.; Maeda, H.; Oishi, S.; Tanaka, T.; Fujii, N.; Ohno, H. Synthesis of fused and linked bicyclic nitrogen heterocycles by palladium-catalyzed domino cyclization of propargyl bromides. *Chem. Eur. J.* **2010**, 16(28) 8410-8418.
- 61 Hirano, K.; Inaba, Y.; Watanabe, T.; Oishi, S.; Fujii, N.; Ohno, H. Gold-catalyzed intramolecular alkyne cycloisomerization cascade: direct synthesis of aryl-annulated[*a*]carbazoles from aniline-substituted diethynylarenes. *Adv. Syn. Catalyst.* **2010**, 352(2) 368-372.
- 62 Nishizawa, K.; Nishiyama, H.; Oishi, S.; Tanahara, N.; Kotani, H.; Mikami, Y.; Toda, Y.; Evans, B. J.; Peiper, S. C.; Saito, R.; Watanabe, J.; Fujii, N.; Ogawa, O. Fluorescent imaging of high grade bladder cancer using a specific antagonist for chemokine receptor CXCR4. *Int. J. Cancer*, **2010**, 127(5) 1180-1187.
- 63 Mizuhara, T.; Oishi, S.; Fujii, N.; Ohno, H. Efficient synthesis of pyrimido[1,2-c][1,3]benzothiazin-6-imines and related tricyclic heterocycles by  $S_NAr$ -type C-S, C-N, or C-O bond formation with heterocumulenes. *J. Org. Chem.* **2010**, 75(1) 265-268.
- 64 Inuki, S.; Yoshimitsu, Y.; Oishi, S.; Fujii, N.; Ohno, H. Ring-construction/stereoselective functionalization cascade: total synthesis of pachastrissamine (jaspine B) through palladium-catalyzed bis-cyclization of bromoallenes. *Org. Lett.* **2009**, 11(19) 4478-4481.
- 65 Ohta, Y.; Chiba, H.; Oishi, S.; Fujii, N.; Ohno, H. Construction of nitrogen heterocycles bearing an aminomethyl group by copper-catalyzed domino three-component coupling-cyclization. *J. Org. Chem.* **2009**, 74(18) 7052-7058.

- 66 Ohta, Y.; Kubota, Y.; Watabe, T.; Chiba, H.; Oishi, S.; Fujii, N.; Ohno, H. Rapid access to 3-(aminomethyl)isoquinoline-fused polycyclic compounds by copper-catalyzed four-component coupling, cascade cyclization, and oxidation. *J. Org. Chem.* **2009**, 74(16) 6299-6302.
- 67 Tanaka, T.; Nomura, W.; Narumi, T.; Esaka, A.; Oishi, S.; Ohashi, N.; Itotani, K.; Evans, B. J.; Wang, Z.; Peiper, S. C.; Fujii, N.; Tamamura, H. Structure-activity relationship study on artificial CXCR4 ligands possessing the cyclic pentapeptide scaffold: the exploration of amino acid residues of pentapeptides by substitutions of several aromatic amino acids. *Org. Biomol. Chem.* **2009**, 7(18) 3805-3809.
- 68 Evans, B.J.; Wang, Z.; Broach, J.R.; Oishi, S.; Fujii, N.; Peiper S.C. Expression of CXCR4, a G-protein-coupled receptor for CXCL12 in yeast identification of new-generation inverse agonists. *Methods Enzymol.* **2009**, 460, 399-412.
- 69 Kobayashi, K.; Narumi, T.; Oishi, S.; Ohno, H.; Fujii, N. Amino acid-based synthesis of trifluoromethylalkene dipeptide isosteres by alcohol-assisted nucleophilic trifluoromethylation and organozinc-copper-mediated S<sub>N</sub>2' alkylation. *J. Org. Chem.* **2009**, 74(12) 4626-4629.
- 70 Suzuki, Y.; Ohta, Y.; Oishi, S.; Fujii, N.; Ohno, H. Efficient synthesis of aminomethylated pyrroloindoles and dipyrrolopyridines via controlled copper-catalyzed domino multi-component coupling and bis-cyclization. *J. Org. Chem.* **2009**, 74(11) 4246-4251.
- 71 Watanabe, T.; Oishi, S.; Fujii, N.; Ohno, H. Palladium-catalyzed direct synthesis of carbazoles via one-pot *N*-arylation and oxidative biaryl coupling: synthesis and mechanistic study. *J. Org. Chem.* **2009**, 74(13) 4720-4726.
- 72 Mizuhara, T.; Inuki, S.; Oishi, S.; Fujii, N.; Ohno, H. Cu(II)-mediated oxidative intermolecular ortho C-H functionalisation using tetrahydropyrimidine as the directing group. *Chem. Commun.* **2009**, (23) 3413-3415.
- 73 Ohta, Y.; Oishi, S.; Fujii, N.; Ohno, H. Facile synthesis of 1,2,3,4-tetrahydro- $\beta$ -carbolines by one-pot domino three-component indole formation and nucleophilic cyclization. *Org. Lett.* **2009**, 11(9) 1979-1982.
- 74 Ueno, M.; Kodama, E.N.; Shimura, K.; Sakurai, Y.; Kajiwara, K.; Sakagami, Y.; Oishi, S.; Fujii, N.; Matsuoka, M. Synonymous mutations in stem-loop III of Rev responsive elements enhance HIV-1 replication impaired by primary mutations for resistance to enfuvirtide. *Antiviral Res.* **2009**, 82(1) 67-72.
- 75 Katagiri, F.; Nagai, K.; Kida, A.; Tomita, K.; Oishi, S.; Takeyama, M.; Doi, R.; Fujii, N. Clinical significance of plasma metastin level in pancreatic cancer patients. *Oncol. Rep.* **2009**, 21, 815-819.
- 76 Nagai, K.; Doi, R.; Katagiri, F.; Ito, T.; Kida, A.; Koizumi, M.; Masui, T.; Kawaguchi, Y.; Tomita, K.; Oishi, S.; Fujii, N.; Uemoto, S. Prognostic value of metastin expression in human pancreatic cancer. *J. Exp. Clin. Cancer Res.* **2009**, 28, 9.
- 77 Naito, T.; Izumi, K.; Kodama, E.; Sakagami, Y.; Kajiwara, K.; Nishikawa, H.; Watanabe, K.; Sarafianos, S.G.; Oishi, S.; Fujii, N.; Matsuoka, M. SC29EK, a peptide fusion inhibitor with enhanced  $\alpha$ -helicity, inhibits replication of human immunodeficiency virus type 1 mutants resistant to enfuvirtide. *Antimicrob. Agents Chemother.* **2009**, 53(3) 1013-1018.
- 78 Izumi, K.; Kodama, E.; Shimura, K.; Sakagami, Y.; Watanabe, K.; Ito, S.; Watabe, T.; Terakawa, Y.; Nishikawa, H.; Sarafianos, S. G.; Kitaura, K.; Oishi, S.; Fujii, N.; Matsuoka, M. Design of peptide-based inhibitors of HIV-1 strains resistant to T-20. *J. Biol. Chem.* **2009**, 284(8) 4914-4920.
- 79 Kitaori, T.; Ito, H.; Schwarz, E.M.; Tsutsumi, R.; Yoshitomi, H.; Oishi, S.; Nakano, M.; Fujii, N.; Nagasawa, T.; Nakamura, T. SDF-1/CXCR4 signaling is critical for the recruitment of mesenchymal stem cells to the fracture site during skeletal repair. *Arthritis Rheum.* **2009**, 60(3) 813-823.
- 80 Mizukoshi, F.; Baba, K.; Goto-Koshino, Y.; Setoguchi-Mukai, A.; Fujino, Y.; Ohno, K.; Tamamura, H.; Oishi, S.; Fujii, N.; Tsujimoto, H. Inhibitory effect of newly developed CXC-chemokine receptor 4 antagonists on the infection with feline immunodeficiency virus. *Vet. Microbiol.* **2009**, 136(1-2) 155-159.
- 81 Mizukoshi, F.; Baba, K.; Goto, Y.; Setoguchi, A.; Fujino, Y.; Ohno, K.; Oishi, S.; Kodera, Y.; Fujii, N.; Tsujimoto, H. Antiviral activity of membrane fusion inhibitors that target gp40 of the feline immunodeficiency virus envelope protein. *J. Vet. Med. Sci.* **2009**, 71(1) 121-124.
- 82 Nishikawa, H.; Nakamura, S.; Kodama, E.; Ito, S.; Kajiwara, K.; Izumi, K.; Sakagami, Y.; Oishi, S.; Ohkubo, T.; Kobayashi, Y.; Otaka, A.; Fujii, N.; Matsuoka, M. Electrostatically constrained  $\alpha$ -helical peptide inhibitors replication of HIV-1 resistant to enfuvirtide. *Int. J. Biochem. Cell Biol.* **2009**, 41(4) 891-899.

- 83 Inuki, S.; Oishi, S.; Fujii, N.; Ohno, H. Total synthesis of ( $\pm$ )-lysergic acid, lysergol, and isolysergol by palladium-catalyzed domino cyclization of amino allenes bearing a bromoindolyl group. *Org. Lett.* **2008**, 10(22) 5239-5242.
- 84 Mikami, S.; Nakase, H.; Yamamoto, S.; Takeda, Y.; Yoshino, T.; Kasahara, K.; Ueno, S.; Uza, N.; Oishi, S.; Fujii, N.; Nagasawa, T.; Chiba, T. Blockade of CXCL12/CXCR4 axis ameliorates murine experimental colitis. *J. Pharmacol. Exp. Ther.* **2008**, 327(2) 383-392.
- 85 Ohta, Y.; Chiba, H.; Oishi, S.; Fujii, N.; Ohno, H. Concise synthesis of indole-fused 1,4-diazepines through copper(I)-catalyzed domino three-component coupling-cyclization-N-arylation under microwave irradiation. *Org. Lett.* **2008**, 10(16) 3535-3538.
- 86 Yano, Y.; Yano, A.; Oishi, S.; Sugimoto, Y.; Tsujimoto, G.; Fujii, N.; Matsuzaki, K. Coiled-coil tag-robe system for quick labeling of membrane receptors in living cells. *ACS Chem. Biol.* **2008**, 3(6) 341-345.
- 87 Okano, A.; Mizutani, T.; Oishi, S.; Tanaka, T.; Ohno, H.; Fujii, N. Palladium-catalysed biscyclisation of allenic bromoalkenes through zipper-mode cascade. *Chem. Commun.* **2008**, (30) 3534-3536.
- 88 Nishikawa, H.; Kodama, E.; Sakakibara, A.; Fukudome, A.; Izumi, K.; Oishi, S.; Fujii, N.; Matsuoka, M. Novel screening systems for HIV-1 fusion mediated by two extra-virion heptad repeats of gp41. *Antivir. Res.* **2008**, 80(1) 71-76.
- 89 Watanabe, T.; Oishi, S.; Fujii, N.; Ohno, H. Palladium-catalyzed  $sp^3$  C-H activation of simple alkyl groups: direct preparation of indoline derivatives from *N*-alkyl-2-bromoanilines. *Org. Lett.* **2008**, 10(9) 1759-1762.
- 90 Inokuchi, E.; Narumi, T.; Niida, A.; Tomita, K.; Oishi, S.; Ohno, H.; Fujii, N. Efficient synthesis of trifluoromethyl and related trisubstituted alkene dipeptide isosteres by palladium-catalyzed carbonylation of amino acid-derived allylic carbonates. *J. Org. Chem.* **2008**, 73(10) 3942-3945.
- 91 Suzuki, Y.; Cluzeau, J.; Hara, T.; Hirasawa, A.; Tsujimoto, G.; Oishi, S.; Ohno, H.; Fujii, N. Structure-activity relationship of pyrazine-based CK2 inhibitors: synthesis and evaluation of 2,6-disubstituted pyrazines and 4,6-disubstituted pyrimidines. *Arch. Pharm.* **2008**, 341(9) 554-561.
- 92 Narumi, T.; Tomita, K.; Inokuchi, E.; Kobayashi, K.; Oishi, S.; Ohno, H.; Fujii, N. Diastereoselective synthesis of highly functionalized fluoroalkene dipeptide isosteres and its application to Fmoc-based solid phase synthesis of a cyclic pentapeptide mimetic. *Tetrahedron*, **2008**, 64(19) 4332-4346.
- 93 Ohta, Y.; Oishi, S.; Fujii, N.; Ohno, H. Facile synthesis of 3-(aminomethyl)isoquinolines by copper-catalysed domino four-component coupling and cyclisation. *Chem. Commun.* **2008**, (7) 835-837.
- 94 Watanabe, T.; Oishi, S.; Fujii, N.; Ohno, H. Gold-catalyzed hydroarylation of allenes: a highly regioselective carbon-carbon bond formation producing six-membered rings. *Org. Lett.* **2007**, 9(23) 4821-4824.
- 95 Watanabe, T.; Ueda, S.; Inuki, S.; Oishi, S.; Fujii, N.; Ohno, H. One-pot synthesis of carbazoles by palladium-catalyzed N-arylation and oxidative coupling. *Chem. Commun.* **2007**, (43) 4516-4518.
- 96 Tamamura, H.; Tanaka, T.; Tsutsumi, H.; Nemoto, K.; Mizokami, S.; Ohashi, N.; Oishi, S.; Fujii, N. Versatile use of acid-catalyzed ring-opening of  $\beta$ -aziridinyl- $\alpha,\beta$ -enoates to stereoselective synthesis of peptidomimetics. *Tetrahedron*, **2007**, 63(37) 9243-9254.
- 97 Narumi, T.; Tomita, K.; Inokuchi, E.; Kobayashi, K.; Oishi, S.; Ohno, H.; Fujii, N. Facile synthesis of fluoroalkenes by palladium-catalyzed reductive defluorination of allylic gem-difluorides. *Org. Lett.* **2007**, 9(17) 3465-3468.
- 98 Ogo, N.; Oishi, S.; Matsuno, K.; Sawada, J.; Fujii, N.; Asai, A. Synthesis and biological evaluation of L-cysteine derivatives as mitotic kinesin Eg5 inhibitors. *Bioorg. Med. Chem. Lett.* **2007**, 17(14), 3921-3924.
- 99 Katagiri, F.; Tomita, K.; Oishi, S.; Takeyama, M.; Fujii, N. Establishment and clinical application of enzyme immunoassays for determination of luteinizing hormone releasing hormone and metastatin. *J. Pept. Sci.* **2007**, 13(6) 422-429.
- 100 Kang, S.-U.; Choi, W. J.; Oishi, S.; Lee, K.; Karki, R. G.; Worthy, K. M.; Bindu, L. K.; Nicklaus, M. C.; Fisher, R. J.; Burke, T. R., Jr. Examination of acylated 4-aminopiperidine-4-carboxylic acid residues in the pTyr+1 position of Grb2 SH2 domain-binding tripeptides. *J. Med. Chem.* **2007**, 50(8) 1978-1982.
- 101 Ohno, H.; Ohta, Y.; Oishi, S.; Fujii, N. Direct synthesis of 2-(aminomethyl)indoles through copper(I)-catalyzed domino three-component coupling and cyclization reactions. *Angew. Chem. Int. Ed.* **2007**, 46(13) 2295-2298.

- 102 Kato-Takagaki, K.; Suzuki, N.; Yokoyama, F.; Takaki, D.; Umezawa, K.; Higo, J.; Mochizuki, M.; Kikkawa, Y.; Oishi, S.; Utani, A.; Nomizu, M. Cyclic peptide analysis of the biologically active loop region in the laminin R3 chain LG4 module demonstrates the importance of peptide conformation on biological activity. *Biochemistry*, **2007**, 46(7) 1952-1960.
- 103 Narumi, T.; Niida, A.; Tomita, K.; Oishi, S.; Otaka, A.; Ohno, H.; Fujii, N. A novel one-pot reaction involving organocopper-mediated reduction/transmetalation/asymmetric alkylation, leading to the diastereoselective synthesis of functionalized (*Z*)-fluoroalkene dipeptide isosteres. *Chem. Commun.* **2006**, (45) 4720-4722.
- 104 Baba, M.; Hong, S.-B.; Sharma, N.; Warren, M.B.; Nickerson, M.L.; Iwamatsu, A.; Esposito, D.; Gillette, W.K.; Hopkins, R.F.; III; Hartley, J.L.; Furihata, M.; Oishi, S.; Zhen, W.; Burke, T.R., Jr.; Linehan, W.M.; Schmidt, L.S.; Zbar, B. Folliculin encoded by the *BHD* gene interacts with a novel binding protein, FNIP1, and AMPK, and is involved in AMPK and mTOR signaling. *Proc. Natl. Acad. Sci.* **2006**, 103(42) 15552-15557.
- 105 Niida, A.; Mizumoto, M.; Narumi, T.; Inokuchi, E.; Oishi, S.; Ohno, H.; Otaka, A.; Kitaura, K.; Fujii, N. Synthesis of (*Z*)-alkene and (*E*)-fluoroalkene containing diketopiperazine mimetics utilizing organocopper-mediated reduction-alkylation and diastereoselectivity examination using DFT calculations. *J. Org. Chem.* **2006**, 71(11) 4118-4129.
- 106 Niida, A.; Tanigaki, H.; Inokuchi, E.; Sasaki, Y.; Oishi, S.; Ohno, H.; Tamamura, H.; Wang, Z.; Peiper, S. C.; Kitaura, K.; Otaka, A.; Fujii, N. Stereoselective synthesis of 3,6-disubstituted-3,6-dihydropyridin-2-ones as potential diketopiperazine mimetics using organocopper-mediated anti- $S_N2'$  reactions and their use in the preparation of low molecule CXCR4 antagonists. *J. Org. Chem.* **2006**, 71(10) 3942-3951.
- 107 Niida, A.; Tomita, K.; Mizumoto, M.; Tanigaki, H.; Terada, T.; Oishi, S.; Otaka, A.; Inui, K.-I.; Fujii, N. Unequivocal synthesis of (*Z*)-alkene and (*E*)-fluoroalkene dipeptide isosteres to probe structural requirements of the peptide transporter PEPT1. *Org. Lett.* **2006**, 8(4) 613-616.
- 108 Niida, A.; Wang, Z.; Tomita, K.; Oishi, S.; Tamamura, H.; Otaka, A.; Navenot, J.-M.; Broach, J. R.; Peiper, S. C.; Fujii, N. Design and synthesis of downsized metastin (45-54) analogs with maintenance of high GPR54 agonistic activity. *Bioorg. Med. Chem. Lett.* **2006**, 16(1) 134-137.
- 109 Niida, A.; Oishi, S.; Sasaki, Y.; Mizumoto, M.; Tamamura, H.; Fujii, N.; Otaka, A. Facile access to (*Z*)-alkene-containing diketopiperazine mimetics utilizing organocopper-mediated anti- $S_N2'$  reactions. *Tetrahedron Lett.* **2005**, 46(24) 4183-4186.
- 110 Tamamura, H.; Araki, T.; Ueda, S.; Wang, Z.; Oishi, S.; Esaka, A.; Trent, J. O.; Nakashima, H.; Yamamoto, N.; Peiper, S. C.; Otaka, A.; Fujii, N. Identification of novel low molecular weight CXCR4 antagonists by structural tuning of cyclic tetrapeptide scaffolds. *J. Med. Chem.* **2005**, 48(9) 3280-3289.
- 111 Ichikawa, N.; Kasai, S.; Suzuki, N.; Nishi, N.; Oishi, S.; Fujii, N.; Kadoya, Y.; Hatori, K.; Mizuno, Y.; Nomizu, M.; Arikawa-Hirasawa, E. Identification of neurite outgrowth active sites on the laminin  $\alpha$ 4 chain G domain. *Biochemistry*, **2005**, 44(15) 5755-5762.
- 112 Shi, Z.-D.; Karki, R. G.; Oishi, S.; Worthy, K. M.; Bindu, L. K. Dharmawardana, P. G.; Nicklaus, M. C.; Bottaro, D. P. Fisher, R. J.; Burke, T. R., Jr. Utilization of a nitrobenzoxadiazole (NBD) fluorophore in the design of a Grb2 SH2 domain-binding peptide mimetic. *Bioorg. Med. Chem. Lett.* **2005**, 15(5) 1385-1388.
- 113 Yokoyama, F.; Suzuki, N.; Haruki, M.; Nishi, N.; Oishi, S.; Fujii, N.; Utani, A.; Kleinman, H. K.; Nomizu, M. Cyclic Peptides from the Loop Region of the Laminin  $\alpha$ 4 Chain LG4 Module Show Enhanced Biological Activity over Linear Peptides. *Biochemistry*, **2004**, 43(42) 13590-13597.
- 114 Otaka, A.; Watanabe, J.; Yukimasa, A.; Sasaki, Y.; Watanabe, H.; Kinoshita, T.; Oishi, S.; Tamamura, H.; Fujii, N. SmI<sub>2</sub>-mediated reduction of  $\gamma,\gamma$ -difluoro- $\alpha,\beta$ -enoates with application to the synthesis of functionalized (*Z*)-fluoroalkene-type dipeptide isosteres. *J. Org. Chem.* **2004**, 69(5) 1634-1645.
- 115 Masui, T.; Doi, R.; Mori, T.; Toyoda, E.; Koizumi, M.; Kami, K.; Ito, D.; Peiper, S. C.; Broach, J. R.; Oishi, S.; Niida, A.; Fujii, N.; Imamura, M. Metastin and its variant forms suppress migration of pancreatic cancer cells. *Biochem. Biophys. Res. Commun.* **2004**, 315(1) 85-92.
- 116 Suzuki, N.; Nakatsuka, H.; Mochizuki, M.; Nishi, N.; Kadoya, Y.; Utani, A.; Oishi, S.; Fujii, N.; Kleinman, H. K.; Nomizu, M. Biological activities of homologous loop regions in the laminin  $\alpha$  chain G domains. *J. Biol. Chem.* **2003**, 278(46) 45697-45705.

- 117 Otaka, A.; Yukimasa, A.; Watanabe, J.; Sasaki, Y.; Oishi, S.; Tamamura, H.; Fujii, N. Application of samarium diiodide ( $\text{SmI}_2$ )-induced reduction of  $\gamma$ -acetoxy- $\alpha,\beta$ -enoates with  $\alpha$ -specific kinetic electrophilic trapping for the synthesis of amino acid derivatives. *Chem. Commun.* **2003**, (15) 1834-1835.
- 118 Ogawa, M.; Hatano, K.; Oishi, S.; Kawasumi, Y.; Fujii, N.; Kawaguchi, M.; Doi, R.; Imamura, M.; Yamamoto, M.; Ajito, K.; Mukai, T.; Saji, H.; Ito, K. Direct electrophilic radiofluorination of a cyclic RGD peptide for in vivo  $\alpha_v\beta_3$  integrin related tumor imaging. *Nucl. Med. Biol.* **2003**, 30(1) 1-9.
- 119 Otaka, A.; Katagiri, F.; Kinoshita, T.; Oishi, S.; Tamamura, H.; Fujii, N. Regio- and stereoselective synthesis of (*E*)-alkene trans-Xaa-Pro dipeptide mimetics utilizing organocopper-mediated anti- $\text{S}_{\text{N}}2'$  Reactions. *J. Org. Chem.* **2002**, 67(17) 6152-6161.
- 120 Ohno, H.; Miyamura, K.; Tanaka, T.; Oishi, S.; Toda, A.; Takemoto, Y.; Fujii, N.; Ibuka, T. Synthesis of allenes from allylic alcohol derivatives bearing a bromine atom using a palladium(0)/diethylzinc system. *J. Org. Chem.* **2002**, 67(4) 1359-1367.
- 121 Tamamura, H.; Omagari, A.; Hiramatsu, K.; Oishi, S.; Habashita, H.; Kanamoto, T.; Gotoh, K.; Yamamoto, N.; Nakashima, H.; Otaka, A.; Fujii, N. Certification of the critical importance of L-3-(2-naphthyl)alanine at position 3 of a specific CXCR4 inhibitor, T140, leads to an exploratory performance of its downsizing study, *Bioorg. Med. Chem.* **2002**, 10(5) 1417-1426.
- 122 Tamamura, H.; Hiramatsu K.; Miyamoto K.; Omagari A.; Oishi S.; Nakashima, H.; Yamamoto, N.; Kuroda, Y.; Nakagawa, T.; Otaka, A.; Fujii, N. Synthesis and evaluation of pseudopeptide analogues of a specific CXCR4 inhibitor, T140: The insertion of an (*E*)-alkene dipeptide isostere into the  $\beta\text{II}'$ -turn moiety. *Bioorg. Med. Chem. Lett.* **2002**, 12(6) 923-928.
- 123 Kawaguchi, M.; Hosotani, R.; Ohishi, S.; Fujii, N.; Tulachan, S. S.; Koizumi, M.; Toyoda, E.; Masui, T.; Nakajima, S.; Tsuji, S.; Ida, J.; Fujimoto, K.; Wada, M.; Doi, R.; Imamura, M. A novel synthetic Arg-Gly-Asp-containing peptide cyclo(-RGDF=V-) is the potent inhibitor of angiogenesis. *Biochem. Biophys. Res. Commun.* **2001**, 288(3) 711-717.
- 124 Ohno, H.; Anzai, M.; Toda, A.; Ohishi, S.; Fujii, N.; Tanaka, T.; Takemoto, Y.; Ibuka, T. Stereoselective  $\hookrightarrow$  synthesis of 2-alkenylaziridines and 2-alkenylazetidines by palladium-catalyzed intramolecular amination of  $\alpha$ - and  $\beta$ -amino allenes. *J. Org. Chem.* **2001**, 66(14) 4904-4914.
- 125 Otaka, A.; Watanabe, H.; Yukimasa, A.; Oishi, S.; Tamamura, H.; Fujii, N. New access to  $\alpha$ -substituted (*Z*)-fluoroalkene dipeptide isosteres utilizing organocopper reagents under ‘reduction-oxidative alkylation (R-OA)’ conditions. *Tetrahedron Lett.* **2001**, 42(32) 5443-5446.
- 126 Tamamura, H.; Sugioka, M.; Odagaki, Y.; Omagari, A.; Kan, Y.; Oishi, S.; Nakashima, H.; Yamamoto, N.; Peiper, S. C.; Hamanaka, N.; Otaka, A.; Fujii, N. Conformational study of a highly specific CXCR4 inhibitor, T140, disclosing the close proximity of its intrinsic pharmacophores associated with strong anti-HIV activity. *Bioorg. Med. Chem. Lett.* **2001**, 11(3) 359-362.
- 127 Ohno, H.; Toda, A.; Oishi, S.; Tanaka, T.; Takemoto, Y.; Fujii, N.; Ibuka, T. Novel synthesis of chiral terminal allenes via palladium(0)-catalyzed reduction of mesylates of 2-bromoalk-2-en-1-ols bearing a protected amino group, using diethylzinc. *Tetrahedron Lett.* **2000**, 41(26) 5131-5134.
- 128 Tamamura, H.; Omagari, A.; Oishi, S.; Kanamoto, T; Yamamoto, N.; Peiper, S. C.; Nakashima, H.; Otaka, A.; Fujii, N. Pharmacophore identification of a specific CXCR4 inhibitor, T140, leads to development of effective anti-HIV agents with very high selectivity indexes. *Bioorg. Med. Chem. Lett.* **2000**, 10(23) 2633-2637.
- 129 Buffet, P.; Gamain, B.; Scheidig, C.; Baruch, D.; Smith, J.; Hernandez-Rivas, R.; Pouvelle, B.; Oishi, S.; Fujii, N.; Fusai, T.; Parzy, D.; Miller, L.; Gysin, J.; Scherf, A. *Plasmodium falciparum* domain mediating adhesion to chondroitin sulfate A: a receptor for human placental infection. *Proc. Natl. Acad. Sci.* **1999**, 96(22) 12743-12748.

## 総説・著書

- 1 Popiel, H. A.; Burke, J. R.; Strittmatter, W. J.; Oishi, S.; Fujii, N.; Toda, T.; Wada K.; Nagai, Y. The aggregation inhibitor peptide QBP1 as a therapeutic molecule for the polyglutamine neurodegenerative diseases. *J. Amino Acids*, **2011**, 2011, 265084.
- 2 大石真也、藤井信孝 ケモカイン受容体 CXCR4 とそのアンタゴニスト 和光純薬時報 **2010**, 78(1) 2-5.
- 3 Hyunsuk, S.; Oishi, S.; Fujii, N. Chemokine receptor CXCR4 as a therapeutic target for neuroectodermal tumors. *Semin. Cancer Biol.* **2009**, 19(2) 123-134.
- 4 Oishi, S.; Narumi, T.; Ohno, H.; Otaka, A.; Fujii, N. Synthesis of highly functionalized alkene dipeptide isosteres and its application to the structure-activity relationship study on bioactive peptides. *J. Synth. Org. Chem. Jpn.* **2008**, 66(9), 846-857.
- 5 掛谷秀昭、大石真也、矢倉徹、藤井信孝 化合物ライブラリー インシリコ創薬科学 (3.2 節・京都廣川書店) **2008**, 46-63.
- 6 大石真也、藤井信孝 ハイスループットスクリーニング インシリコ創薬科学 (3.3 節・京都廣川書店) **2008**, 63-74.
- 7 大石真也、藤井信孝 ペプチドをリードとした創薬 蛋白質 核酸 酵素 増刊 ケミカルバイオロジー 長野哲雄・長田裕之・菊地和也・上杉志成編集, **2007**, 52(13), 1696-1701.
- 8 大石真也、藤井信孝 ケミカルバイオロジーを基盤とするペプチド創薬 遺伝子医学 MOOK8 ペプチドと創薬 寒川賢治・南野直人編集, **2007**, 87-91.
- 9 Tsutsumi, H.; Tanaka, T.; Ohashi, N.; Masuno, H.; Tamamura, H.; Hiramatsu, K.; Araki, T.; Ueda, S.; Oishi, S.; Fujii, N. The therapeutic potential of the chemokine receptor CXCR4 antagonists as multi-functional agents. *Biopolymers*, **2007**, 88(2), 279-289.
- 10 藤井信孝、大石真也、大高章 プロテオミクスを基盤とするケミカルバイオロジー—情報から制御へ バイオニクス **2006**, 3(10), 46-51.
- 11 Oishi, S. Structure-activity relationship studies on cyclic RGD peptides utilizing novel alkene dipeptide isosteres. *Yakugaku Zasshi*, **2004**, 124(5), 269-277.